

Antifungal Reference Document for Adult Patients at Stanford Health Care

Formulations Available at SHC		Therapeutic Drug Monitoring	Dose Adjustments	Clinical Pearls	
Voriconazole	Tablets (50 mg, 200 mg)	Administer on an empty stomach*	<p><u>When to consider TDM</u></p> <p>Recommended for all patients receiving voriconazole given inter- and intra-patient variability and exposure-response and toxicity profile</p> <p><u>Type of level</u> Trough (i.e. prior to the next dose or ~10-12 hours after the last dose)</p> <p><u>When to get a level</u> Day ≥ 4</p> <p><u>Prophylaxis</u> ≥1 to ≤ 5.5 mg/L</p> <p><u>Treatment</u> ≥ 1 to ≤ 5.5 mg/L (may consider > 2 mg/L in certain clinical scenarios in consultation with ID)</p> <p><u>When to consider additional levels</u></p> <ul style="list-style-type: none"> • After a change in dose • Introduction or discontinuation of drugs with significant interaction potential • Severe diarrhea and receiving oral formulations • Disease progression • Concern for non-adherence • Concern for toxicity 	<p><u>Trough level < 1 mg/L:</u> Non-linear pharmacokinetics: dose adjustments do not lead to proportional changes in serum levels. Increase daily dose by 50 – 100 mg & re-check a level in 4-7 days</p> <p><u>Trough level > 5.5 mg/L:</u> Decrease daily dose by 50 – 100 mg & re-check a level in 4-7 days. May need to consider holding dose(s) in certain clinical scenarios</p> <p>Discuss with ID consult service or clinical pharmacist as needed.</p>	<p><u>PK/PD</u> Enteral formulations have high bioavailability (especially on an empty stomach); consider tablets or suspension for patients unless contraindicated.</p> <p>Serum levels > 5.5 mg/L are associated with visual hallucinations/risk of neurotoxicity and may be associated with hepatotoxicity</p> <p>Significant interpatient variability in serum levels due to polymorphisms in <i>CYP2C19</i></p> <p>Use adjusted body weight in setting of obesity‡</p> <p><u>Drug-Drug Interactions</u> Clinically significant drug-drug interactions.</p> <p><u>Adverse Drug Reactions</u> Use has been associated with transient visual disturbances, QTc prolongation, transaminase abnormalities, and, with long-term use, fluorosis and dermatological complications, including cutaneous malignancies</p>
	Oral suspension (40 mg/mL)				
	Intravenous	Formulated in sulfobutylether-β-cyclodextrin. Clinical significance of accumulation in renal insufficiency is debated. Toxicity has only been demonstrated in animal models and at doses far above normal human doses. Weigh risk vs benefit and consider enteral therapy when appropriate			

*Administer 1 hour before or 1 hour after meals

‡Refer to [Antimicrobial Dosing Reference Guide](#) for dosing recommendations

Posaconazole	Formulations Available at SHC		Therapeutic Drug Monitoring	Dose Adjustments	Clinical Pearls
	Delayed Release tablet (DRT) (100 mg)	Administer with food.	<p>When to consider TDM Recommended for patients receiving posaconazole for treatment.</p> <p>For prophylaxis, TDM may not be necessary if receiving DRT/IV formulations. Consider TDM on a case-by-case basis (e.g. extremes of body weight, concern for breakthrough infection, crushing DRTs etc.), as per protocol, or in consultation with ID</p> <p>Type of level ‘Trough’ (random level acceptable)</p> <p>When to get a level* Day 5 – 7</p> <p>Prophylaxis ≥ 700 ng/mL**</p> <p>Treatment ≥ 1,000 ng/mL (≥ 1,250 ng/mL may be considered in certain clinical scenarios in consultation with ID)</p> <p>When to consider additional levels</p> <ul style="list-style-type: none"> • After a change in dose • Morbid obesity • Introduction or discontinuation of drugs with significant interaction potential • Diarrhea and receiving oral formulations • Disease progression • Concern for non-adherence • Concern for toxicity 	<p>Trough level below target and on DR tablet/intravenous formulation:</p> <p>Assume linear/proportional pharmacokinetics. Increase dose in 100 mg increments if near target. In patients with very low levels (e.g. < 500 ng/mL), larger dose adjustments and more frequent dosing intervals may be necessary (e.g., q12hr dosing). Re-check a level in 5 – 7 days</p> <p>Discuss with ID consult service or clinical pharmacist as needed.</p> <p>No toxicity threshold has been established. Limited data suggest an association with higher serum posaconazole concentrations and pseudohypoaldosteronism. Some licensing authorities suggest an upper end of 3,750 ng/mL be considered</p>	<p>PK/PD DRTs have high bioavailability (especially when administered with food) and should be considered for patients unless contraindicated***</p> <p>Posaconazole suspension (non-formulary) is not interchangeable with the DRT/IV preparations due to saturable absorption & erratic pharmacokinetics.</p> <p>For patients without central access & NG tube in place, consider alternative antifungal options or crushing DRTs. There is limited data to support crushing the posaconazole DRT for administration via feeding tubes. If this is done, TDM should be performed</p> <p>Drug-Drug Interactions Clinically significant drug-drug interactions.</p> <p>Adverse Drug Reactions Use has been associated with GI disturbances, QTc prolongation, transaminase abnormalities, and metabolic side effects</p>
Intravenous (IV)	Administer via a central line. Formulated in sulfobutylether-β-cyclodextrin. Clinical significance of accumulation in renal insufficiency is debated. Toxicity has only been demonstrated in animal models and at doses far above normal human doses. Weigh risk vs benefit and consider enteral therapy when appropriate				

* Serum levels obtained at day three may be considered as a surrogate for steady-state levels on a case-by-case basis

**Based on a pooled analysis of PK data from two Phase III clinical trials. Some studies and consensus guidelines have suggested that a lower target of > 500 ng/mL may be acceptable

***Refer to [Pharmacist-Managed Intravenous to Oral Therapy Interchange Protocol](#)

‡Refer to [Antimicrobial Dosing Reference Guide](#) for dosing recommendations

Itraconazole	Formulations Available at SHC		Therapeutic Drug Monitoring	Dose Adjustments*	Clinical Pearls
	Capsule (100 mg)	Administer with a full meal and/or acidic beverage Acid suppression (e.g. PPI or H2RA) should be avoided	<p><u>When to consider TDM</u> Recommended for all patients</p> <p><u>Type of level</u> "Trough" (random level acceptable)</p> <p>HPLC assay measures itraconazole and hydroxyitraconazole levels. <u>Both values should be added to evaluate true level.</u></p>	<p>Random level < 0.5 mg/L: Increase total daily dose by 100 – 200 mg</p> <p>Random level > 5 mg/L: Consider dose reduction if the patient is experiencing an adverse event or transition to an alternative antifungal if clinically appropriate</p>	<p><u>PK/PD</u> Oral solution has higher bioavailability relative to the capsule but may not be well tolerated due to GI side effects. Given differences in relative bioavailability, these products are not directly interchangeable</p> <p>Some centers measure itraconazole levels by bioassay; these levels are not interchangeable with HPLC</p> <p>A new formulation of itraconazole (Tolsura®) with enhanced oral bioavailability was FDA approved in 2018 (not on SHC formulary)</p> <p><u>Drug-Drug Interactions</u> Clinically significant drug-drug interactions.</p> <p><u>Adverse Drug Reactions</u> Use has been associated with GI disturbances, QTc prolongation, heart failure (rare cases of new onset & exacerbation mostly described with IV preparation)</p>
	Oral Solution (10 mg/mL)	Administer on an empty stomach*	<p><u>When to get a level</u> Day 5 – 7 if loading dose administered Day 10 – 14 if no loading dose administered</p> <p><u>Prophylaxis**</u> >0.5 mg/L – < 5 mg/L</p> <p><u>Treatment</u> >1 mg/L – < 5 mg/L (may consider > 2 mg/L in certain clinical scenarios)</p> <p><u>When to consider additional levels</u></p> <ul style="list-style-type: none"> • After a change in dose • Introduction or discontinuation of drugs with significant interaction potential • Diarrhea and receiving oral formulations • Disease progression • Concern for non-adherence • Concern for toxicity 		

*Administer 1 hour before or 1 hour after meals

**Non-linear pharmacokinetics; dose adjustments do not lead to proportional changes in serum levels

HPLC = high-performance liquid chromatography

Isavuconazole	Formulations Available at SHC		Therapeutic Drug Monitoring	Dose Adjustments	Clinical Pearls
	Capsule (186 mg)	Administer without regards to meals	<p><u>When to consider TDM</u></p> <p>Not routinely recommended. Clinical scenarios necessitating TDM should be discussed with ID and/or ASP (e.g. if capsules are opened for administration via feeding tube, concern for breakthrough fungal infection, etc)</p> <p>No exposure-efficacy or exposure-toxicity thresholds have been established</p>		<p><u>PK/PD</u></p> <p>Capsules have high bioavailability and should be considered for patients unless contraindicated*</p> <p>Isavuconazole PK are linear and dose proportional</p> <p><u>Drug-Drug Interactions</u> Clinically significant drug-drug interactions.</p> <p><u>Adverse Drug Reactions</u> Use has been associated with a shortened QT interval, transaminase abnormalities, GI disturbances, peripheral edema, and hypokalemia</p> <p>Emerging data to suggest that opening capsules and administering via NG tubes leads to comparable exposures to administration of intact capsules. Consider TDM if this is pursued.</p>
	Intravenous (IV)	Infuse with an in-line filter (pore size 0.2 to 1.2 micron)			

*Refer to [Pharmacist-Managed Intravenous to Oral Therapy Interchange Protocol](#)

Flucytosine	Formulations Available at SHC		Therapeutic Drug Monitoring	Dose Adjustments**	Clinical Pearls
	Capsule (250 mg, 500 mg)	To reduce or avoid nausea and vomiting, administer a few capsules at a time over 15 minutes until full dose is taken	<p><u>When to consider TDM</u> Recommended for most patients*</p> <p><u>Type of level*</u> Peak (i.e. 2 hours post-dose)</p>	<p><u>When to get a level</u> Within 72 hours of initiation or after 3 to 5 doses</p> <p><u>Target Peak</u> 25 – 100 mg/L</p> <p>Levels > 100 mg/L increase risk of bone marrow suppression and hepatic dysfunction</p>	<p>Peak level < 25 mg/L: Limited data, consider increasing total daily dose by 50% (use clinical judgment)</p> <p>Peak level > 100 mg/L: Hold dose(s) as needed. If due to renal impairment, decrease dosing frequency</p> <p>Acute changes in renal function should prompt consideration for empiric dose adjustment</p>
Oral Suspension	Not commercially available; may be compounded on an as needed basis (discuss with pharmacy)				

*This is a send out lab that takes several days to result. It is important to time level appropriately so that it is clinically actionable

**Demonstrates linear pharmacokinetics

‡Refer to [Antimicrobial Dosing Reference Guide](#) for dosing recommendations

References

1. Voriconazole. Lexi-Drugs. Hudson, OH: Lexicomp, 2020. <http://online.lexi.com/>. Updated January 13, 2020. Accessed April 19, 2020.
2. Pascual A, Calandra T, Bolay S, Buclin T, Bille J, Marchetti O. Voriconazole therapeutic drug monitoring in patients with invasive mycoses improves efficacy and safety outcomes. *Clin. Infect. Dis.* 46(2),201--211 (2008).
3. Ueda K, Nannya Y, Kumano K et al. Monitoring trough concentration of voriconazole is important to ensure successful antifungal therapy and to avoid hepatic damage in patients with hematological disorders. *Int. J. Hematol.* 89(5), 592--599 (2009)
4. Miyakis S, van Hal SJ, Ray J, Marriott D. Voriconazole concentrations and outcome of invasive fungal infections. *Clin. Microbiol. Infect.* 16(7), 927--933 (2010)
5. Dolton MJ, Ray JE, Chen SC, Ng K, Pont LG, McLachlan AJ. Multicenter study of voriconazole pharmacokinetics and therapeutic drug monitoring. *Antimicrob. Agents Chemother.* 56(9), 4793--4799 (2012).
6. Park WB, Kim NH, Kim KH et al. The effect of therapeutic drug monitoring on safety and efficacy of voriconazole in invasive fungal infections: a randomized controlled trial. *Clin. Infect. Dis.* 55(8), 1080--1087 (2012)
7. Tan K, Brayshaw N, Tomaszewski K, Troke P, Wood N. Investigation of the potential relationships between plasma voriconazole concentrations and visual adverse events or liver function test abnormalities. *J. Clin. Pharmacol.* 46(2), 235--243 (2006)
8. Posaconazole. Lexi-Drugs. Hudson, OH: Lexicomp, 2020. <http://online.lexi.com/>. Updated July 16, 2019. Accessed April 19, 2020.
9. Cornely OA, Maertens J, Winston DJ et al. Posaconazole vs fluconazole or itraconazole prophylaxis in patients with neutropenia. *N. Engl. J. Med.* 356(4), 348--359 (2007)
10. Jang SH, Colangelo PM, Gobburu JV. Exposure-response of posaconazole used for prophylaxis against invasive fungal infections: evaluating the need to adjust doses based on drug concentrations in plasma. *Clin. Pharmacol. Ther.* 88(1), 115--119 (2010).
11. Dolton MJ, Ray JE, Chen SC, Ng K, Pont L, McLachlan AJ. Multicenter study of posaconazole therapeutic drug monitoring: exposure-response relationship and factors affecting concentration. *Antimicrob. Agents Chemother.* 56(11), 5503--5510 (2012).
12. Cornely OA, Ullmann AJ. Lack of evidence for exposure-response relationship in the use of posaconazole as prophylaxis against invasive fungal infections. *Clin. Pharmacol. Ther.* 89(3), 351--352 (2011).
13. Prattes, Jürgen, et al. "Posaconazole Plasma Concentrations on Days Three to Five Predict Steady-State Levels." *Antimicrobial Agents and Chemotherapy*, Sept. 2016, pp.5595--99. DOI.org (Crossref), doi:10.1128/AAC.00389-16.
14. Bryant AM, Slain D, Cumpston A, Craig M. A post-marketing evaluation of posaconazole plasma concentrations in neutropenic patients with haematological malignancy receiving posaconazole prophylaxis. *Int J Antimicrob Agents.* 2011 Mar;37(3):266-9. doi: 10.1016/j.ijantimicag.2010.11.021. Epub 2011 Jan 13. PMID: 21236645.
15. Ullmann AJ, Lipton JH, Vesole DH et al. Posaconazole or fluconazole for prophylaxis in severe graft-versus-host disease. *N Engl J Med* 2007; 356: 335--47.
16. Hoeningl M, Raggam RB, Salzer HJ, Valentin T, Valentin A, Zollner-Schwetz I, Strohmaier AT, Seeber K, Wölfler A, Sill H, Krause R. Posaconazole plasma concentrations and invasive mould infections in patients with haematological malignancies. *Int J Antimicrob Agents.* 2012 Jun;39(6):510-3. Doi: 10.1016/j.ijantimicag.2012.02.002. Epub 2012 Apr 3. PMID: 22481057.
17. Walsh TJ, Raad I, Patterson TF, Chandrasekar P, Donowitz GR, Graybill R, Greene RE, Hachem R, Hadley S, Herbrecht R, Langston A, Louie A, Ribaud P, Segal BH, Stevens DA, van Burik JA, White CS, Corcoran G, Gogate J, Krishna G, Pedicone L, Hurdal C, Perfect JR. Treatment of invasive aspergillosis with posaconazole in patients who are refractory to or intolerant of conventional therapy: an externally controlled trial. *Clin Infect Dis.* 2007 Jan 1;44(1):2-12. doi: 10.1086/508774. Epub 2006 Nov 28. PMID: 17143808.
18. Nguyen MH, Davis MR, Wittenberg R, Mchardy I, Baddley JW, Young BY, Odermatt A, Thompson GR. Posaconazole Serum Drug Levels Associated With Pseudohyperaldosteronism. *Clin Infect Dis.* 2020 Jun 10;70(12):2593-2598. doi: 10.1093/cid/ciz741. PMID: 31403165; PMCID: PMC7286374.
19. Itraconazole. Lexi-Drugs. Hudson, OH: Lexicomp, 2020. <http://online.lexi.com/>. Updated August 25, 2019. Accessed April 19, 2020.
20. Dolton, Michael J., and Andrew J. McLachlan. "Optimizing Azole Antifungal Therapy in the Prophylaxis and Treatment of Fungal Infections." *Current Opinion in Infectious Diseases*, Dec. 2014., doi:10.1097/QCO.000000000000103.
21. Denning, David W. "Treatment of Invasive Aspergillosis." *Journal of Infection*, vol. 28, May 1994, doi:10.1016/S0163-4453(94)95941-2.
22. Glasmacher, Axel, et al. "Itraconazole Prevents Invasive Fungal Infections in Neutropenic Patients Treated for Hematologic Malignancies: Evidence From a Meta-Analysis of 3,597 Patients." *Journal of Clinical Oncology*, vol. 21, no. 24, Dec. 2003, doi:10.1200/JCO.2003.04.052.
23. Lestner, Jodi M., et al. "Toxicodynamics of Itraconazole: Implications for Therapeutic Drug Monitoring." *Clinical Infectious Diseases*, Sept. 2009, doi:10.1086/605499.
24. Law, D., et al. "Bioassay for Serum Itraconazole Concentrations Using Hydroxyitraconazole Standards." *Antimicrobial Agents and Chemotherapy*, July 1994, doi:10.1128/AAC.38.7.1561.
25. Isavuconazole. Lexi-Drugs. Hudson, OH: Lexicomp, 2020. <http://online.lexi.com/>. Updated December 12, 2019. Accessed April 19, 2020.
26. Andes, David, et al. "Isavuconazole Concentration in Real-World Practice: Consistency with Results from Clinical Trials." *Antimicrobial Agents and Chemotherapy*, vol. 62, no. 7, May 2018, doi:10.1128/AAC.00585-18.
27. Desai, Amit V., et al. "Exposure-Response Relationships for Isavuconazole in Patients with Invasive Aspergillosis and Other Filamentous Fungi." *Antimicrobial Agents and Chemotherapy*, doi:10.1128/AAC.01034-17.
28. Flucytosine. Lexi-Drugs. Hudson, OH: Lexicomp, 2020. <http://online.lexi.com/>. Updated March 09, 2019. Accessed April 19, 2020.
29. Ashbee HR et al. Therapeutic drug monitoring (TDM) of antifungal agents: guidelines from the British Society for Medical Mycology. *J Antimicrob Chemother.* 2014;69(5):1162-76
30. McCreary EK, Davis MR, Narayanan N, Andes DR, Cattaneo D, Christian R, Lewis RE, Watt KM, Wiederhold NP, Johnson MD. Utility of triazole antifungal therapeutic drug monitoring: Insights from the Society of Infectious Diseases Pharmacists: Endorsed by the Mycoses Study Group Education and Research Consortium. *Pharmacotherapy.* 2023 Oct;43(10):1043-1050. Doi: 10.1002/phar.2850. Epub 2023 Aug 14. PMID: 37459118.
31. Dieringer TD, Schaenman JM, Davis MR. Enteral feeding tube administration with therapeutic drug monitoring of crushed posaconazole tablets and opened isavuconazonium sulfate capsules. *J Antimicrob Chemother.* 2022 Apr 27;77(5):1417-1423. doi: 10.1093/jac/dkac035. PMID: 35174391.
32. McCreary EK, Nguyen MH, Davis MR, Borlagdan J, Shields RK, Anderson AD, Rivosecchi RM, Marini RV, Sacha LM, Silveira FP, Andes DR, Lepak AJ. Achievement of clinical isavuconazole blood concentrations in transplant recipients with isavuconazonium sulphate capsules administered via enteral feeding tube. *J Antimicrob Chemother.* 2020 Oct 1;75(10):3023-3028. Doi: 10.1093/jac/dkaa274. PMID: 32710097; PMCID: PMC7778376.

Document Information:

1. Original Authors/Date: William Alegria, PharmD BCIDP
2. Gatekeeper: Antimicrobial Stewardship Program
3. Review and Renewal Requirement: This document will be reviewed every three years and as required by change of law or practice
4. Approvals: Antimicrobial Subcommittee: 05/28/2020,03/14/2024