

OPERATING ROOM CHECKLIST

Before Procedure (Bedside Nurse)

- Pre-Procedure Checklist Completed in EPIC
- Place infant on a bed with radiant warmer mode (open hotbed or DUAL); set to servo baby temp @ 37°C
- Gather Resuscitation Equipment
 - *Bag/Mask, Oxygen Tanks, Intubation box (if necessary)
- Place transport monitor on bed
- Take Alaris Pump Off Bed
- Set up maintenance IVF on syringe pump
- Have extra IVF syringes available
- Gather warm blankets
- Pick up OR Supplies Packet (NICU Front Desk)
 - **Hat, chemical warmer, Temp Probe, Thermal Reflector, Thermometer, extra clear needleless connector
- Place PIV (max 2 attempts by 2 providers)
- Discuss Line Access with OR Team

IntraOp (OR Staff)

- Plug in bed once returned to OR hallway, power ON
- Set temperature to servo (baby) mode @ 37°C



- Ensure temperature probe is plugged into the bed



PostOp (OR Staff)

- Activate chemical warmer, obtain warm blankets
- Move patient to bed, place directly on chemical warmer. Placed warmed hat.
- Move temp probe to patient



- Verify temperature is set to servo mode to 37°C
- If using an incubator, close the top during transport for optimal thermoregulation.



OPERATING ROOM CHECKLIST

- ✓ Peripheral IV is preferred access; avoid use of central line whenever possible
- ✓ **DO NOT** remove microclaves (clear microclaves have no resistance valves and can run low volumes)
- ✓ Maintain minimum infusion rate in PICC lines at all times

	1.2 Fr	1.9 Fr (single or dual lumen)	2.6 Fr (dual lumen)	3 Fr (single)	4 Fr (single or dual lumen)
Weight	<600 g	>600 g	> 3kg	See footnote*	
Blood draw	No	No	Yes- Red port	Yes	Yes- Single lumen Yes-dual lumen (purple port)
Transfusion	No	No	Yes- Red port	Yes	Yes-Single lumen Yes- dual lumen (purple port)
Minimum rate	1 ml/hr + 0.5U heparin/ml	1 ml/hr + 0.5U heparin/ml	1 ml/hr + 0.5U heparin/ml for <i>each</i> lumen	Heparin lock# (Flush/Discard Volume = 1.5 mL)	Heparin Lock# (Flush/Discard Volume: Single lumen =2 mL Dual lumen = 1.5 mL [white port], 1.5 mL [purple port])

*Note: 3Fr and 4Fr catheters are placed based on ultrasound assessment of vein size, not weight

#Heparin lock is 10 U/mL- minimum dose is 3x the priming volume of the line + add-on devices and is administered every 24h