

# Probes to Monitor Activity of the Paracaspase MALT1

Janna Hachmann,<sup>1,2</sup> Laura E. Edgington-Mitchell,<sup>3,4,8</sup> Marcin Poreba,<sup>7</sup> Laura E. Sanman,<sup>4,5</sup> Marcin Drag,<sup>7</sup> Matthew Bogoy,<sup>4,6</sup> and Guy S. Salvesen<sup>1,\*</sup>

<sup>1</sup>Sanford-Burnham Medical Research Institute, La Jolla, CA 92037, USA

<sup>2</sup>Graduate School of Biomedical Sciences, Sanford-Burnham Medical Research Institute, La Jolla, CA 92037, USA

<sup>3</sup>Cancer Biology Program

<sup>4</sup>Department of Pathology

<sup>5</sup>Department of Chemical and Systems Biology

<sup>6</sup>Department of Microbiology and Immunology

Stanford School of Medicine, Stanford, CA 94305, USA

<sup>7</sup>Division of Bioorganic Chemistry, Wroclaw University of Technology, 50-370 Wroclaw, Poland

<sup>8</sup>Present address: Department of Biochemistry, La Trobe University, Melbourne, VIC 3086, Australia

\*Correspondence: [gsalvesen@sanfordburnham.org](mailto:gsalvesen@sanfordburnham.org)

<http://dx.doi.org/10.1016/j.chembiol.2014.11.011>

## SUMMARY


The human paracaspase mucosa-associated lymphoid tissue lymphoma translocation protein 1 (MALT1) plays a central role in nuclear factor- $\kappa$ B (NF- $\kappa$ B) signaling as both a protease and scaffolding protein. Knocking out MALT1 leads to impaired NF- $\kappa$ B signaling and failure to mount an effective immune response. However, it is unclear to which degree it is the scaffolding function versus the proteolytic activity of MALT1 that is essential. Previous work involving a MALT1 inhibitor with low selectivity suggests that the enzymatic function plays an important role in different cell lines. To help elucidate this proteolytic role of MALT1, we have designed activity-based probes that inhibit its proteolytic activity. The probes selectively label active enzyme and can be used to inhibit MALT1 and trace its activity profile, helping to create a better picture of the significance of the proteolytic function of MALT1.

## INTRODUCTION

The human paracaspase MALT1 (mucosa-associated lymphoid tissue lymphoma translocation protein 1) plays a vital role in nuclear factor- $\kappa$ B (NF- $\kappa$ B) signaling downstream of T and B cell receptors (Lucas et al., 2001; Ruefli-Brasse et al., 2003; Ruland et al., 2003). In addition, other receptors such as Dectin-1 and G-protein-coupled receptors have also been described to signal through MALT1 (Wegener and Krappmann, 2007). To signal, MALT1 forms a complex with the CARD-containing proteins CARMA1 (also known as CARD11) as well as its direct binding partner Bcl10 (Gaide et al., 2002; Pomerantz et al., 2002; Uren et al., 2000; Wang et al., 2002). Once the upstream signal is received, assembly of the CARMA1-Bcl10-MALT1 (CBM) complex ensues, allowing other downstream proteins such as TRAF6 and NEMO (NF- $\kappa$ B essential modulator, also known as IKK- $\gamma$ ) to assemble upon it (Oeckinghaus et al., 2007; Sun et al., 2004; Zhou et al., 2004). From this angle, the presumptive protease MALT1 seems to play an important but nonproteolytic

role. However, upon its discovery almost 15 years ago (Akagi et al., 1999; Dierlamm et al., 1999; Morgan et al., 1999), its homology to the CD clan of proteases was noted (Uren et al., 2000). In 2008, the first proteolytic substrates A20 and Bcl10 were discovered (Coornaert et al., 2008; Rebeaud et al., 2008), followed by CYLD, RelB, Regnase-1, and Roquin (Haifinger et al., 2011; Jeltsch et al., 2014; Staal et al., 2011; Uehata et al., 2013). In addition, the NF- $\kappa$ B-inducing kinase (NIK) (Rosebeck et al., 2011) has been reported to be cleaved by an oncogenic fusion of the cellular inhibitor of apoptosis 2 (cIAP2, also known as AIP2) with MALT1 (cIAP2-MALT1), which occurs with varying frequencies in MALT lymphoma of different sites (Isaacson and Du, 2004). The effects of the cleavages have been described elsewhere (Coornaert et al., 2008; Haifinger et al., 2011; Jeltsch et al., 2014; Rebeaud et al., 2008; Rosebeck et al., 2011; Staal et al., 2011; Uehata et al., 2013), but to summarize, the cleavages of A20 and RelB seem to have mostly fine-tuning effects on NF- $\kappa$ B signaling, whereas the cleavage of Bcl10 is required for increased cellular adhesion to fibronectin after T cell activation, and CYLD cleavage plays a role in JNK signaling. Cleavage of the RNase Regnase-1 and the RNA binding protein Roquin disrupts the regulation of several mRNAs, some of which are involved in NF- $\kappa$ B signaling or are NF- $\kappa$ B target genes. The cleavage of NIK relieves it from intramolecular inhibition through its own TRAF3 domain and allows uncontrolled noncanonical NF- $\kappa$ B signaling. The overall importance of MALT1 for T and B cell development has been shown using MALT1 knockout mice, which fail to mount an effective immune response and NF- $\kappa$ B signaling is prevented (Ruefli-Brasse et al., 2003; Ruland et al., 2003).

Importantly, if the proteolytic activity of MALT1 is inhibited but the protein is still present to fulfill its scaffolding function, NF- $\kappa$ B signaling is decreased but not ablated (Caballar et al., 2013; Coornaert et al., 2008; Lucas et al., 2001; Rebeaud et al., 2008). Thus, questions remain about the importance of the proteolytic activity of MALT1. Initial results using B cell lymphoma cell lines and the broad-spectrum protease inhibitor Z-VPRF-fluoromethylketone (Z-VPRF-FMK) pointed toward an important role in the activated B cell-like subset of diffuse large B cell lymphoma (ABC-DLBCL) (Ferch et al., 2009; Haifinger et al., 2009). When ABC-DLBCL cell lines were treated with Z-VPRF-FMK, cell viability declined, providing proof that inhibiting MALT1 proteolytic activity could be an important treatment option for this



**Figure 1. Synthesis Scheme of the Probes Cy5-LVSR-AOMK and Biotin-LVSR-AOMK**

aggressive form of lymphoma. However, further studies are needed to confirm this, especially considering the low selectivity of the inhibitor used, most likely inhibiting not only MALT1 but also a broad spectrum of other proteases, making clear conclusions difficult. A mouse knock-in model expressing the catalytic mutant of MALT1 (C464A) will certainly provide enlightening insights.

Here we describe inhibitory active site probes of MALT1 that enable visualization and quantification of MALT1 activity at a certain time point or after a stimulus. These activity-based probes consist of a peptide sequence determining enzyme selectivity, an electrophilic warhead, which irreversibly inhibits the targeted protease, and a label that allows detection and activity monitoring. Our work can be read together with a similar independent study in this issue of *Chemistry & Biology* (Eitelhuber et al., 2014).

## RESULTS


### Synthesis of Activity-Based Probes Directed against MALT1

We used the cleavage site specificity determined earlier (LVSR↓) (Hachmann et al., 2012) to design MALT1 activity-based probes. We chose an acyloxymethyl ketone (AOMK) electrophile that preferentially inhibits cysteine proteases and has been used for probe targeting other clan CD proteases (Berger et al., 2006; Edgington et al., 2013; Kato et al., 2005a, 2005b; Powers et al., 2002). We included a Cy5 fluorophore or a biotin label, which

allows affinity purification as well as direct imaging of probe modification (Figure 1).

### Preferential Binding to Activated Recombinant MALT1

To test whether the inhibitors can distinguish between WT and catalytic mutant on the one hand and activated and nonactivated MALT1 on the other hand, we incubated purified recombinant enzyme with Cy5-LVSR-AOMK in the presence or absence of sodium citrate, which we and others have shown previously to activate MALT1 in vitro (Hachmann et al., 2012; Wiesmann et al., 2012). As shown in Figure 2, the activity-based probe can clearly differentiate between MALT1-WT and the catalytic mutant MALT1-C464A. No labeling is seen for the catalytic mutant, whereas Cy5-LVSR-AOMK labels MALT1-WT in a dose-dependent manner, confirming that the probe indeed binds the active site. Labeling is strongly decreased in the absence of sodium citrate, indicating that the probe can distinguish between active and inactive protease. That some binding is seen in non-activating conditions is most likely due to a combination of the high concentration of enzyme present in vitro and substrate-induced (or in this case inhibitor-induced) generation of an active conformation, as previously reported (Wiesmann et al., 2012; Yu et al., 2011). As can be seen in Figure 2A, two protein species are labeled by Cy5-LVSR-AOMK. The upper species represents full-length MALT1, whereas the lower species was determined via mass spectrometry analysis to most likely be a C-terminal fragment encompassing the catalytic domain as well as the



**Figure 2. Cy5-LVSR-AOMK Can Distinguish between MALT1-WT and C464A and Preferentially Binds under Activating Conditions** (A and B) MALT1-WT or MALT1-C464A (400 nM) was incubated with Cy5-LVSR-AOMK at different concentrations in buffer with or without citrate for 30 min at 37°C, followed by TCA precipitation and SDS-PAGE. Fluorescence was scanned (A), followed by staining with a coomassie-based protein stain (B). Data are representative of three independent experiments.

C-terminal extension including the Ig<sub>3</sub> domain (Figure S3 available online). The minimal number of peptides observed in mass spectrometry originating from the N-terminal region of MALT1 is likely carryover in the gel from the full-length protein. The intriguing idea that the C-terminal fragment could be an autoproteolytic cleavage product is contradicted by the fact that the catalytic mutant also shows this product. Since no fragments of this size are seen in cell culture experiments and no convincing cleavage sites are found in the sequence preceding the catalytic domain, we conclude that the cleavage is most likely due to an *E. coli* protease during expression and represents an in vitro artifact leading to two active species.

To confirm that probe binding indeed leads to the inactivation of the enzyme and to obtain evidence that biotin-LVSR-AOMK also binds and inhibits MALT1, we measured the MALT1 rate of inhibition. Pseudo first-order calculations (Table 1) revealed that the rate of inhibition is at least in the range of  $10^3 \text{ M}^{-1}\text{s}^{-1}$  for the inhibitors Cy5-LVSR-AOMK and biotin-LVSR-AOMK, slightly exceeding the rate of inhibition of the previously used inhibitor Z-VRPR-FMK.

### Detection of MALT1 Activity upon Ectopic Expression in Mammalian Cells

We next determined whether the activity-based probes can be used to detect active versus inactive MALT1 in lysates from a

**Table 1.  $k_{\text{obs}}/I$  Values for Inhibition of MALT1 by the Probes/Inhibitors Used in This Study**

Inhibitor	$k_{\text{obs}}/I \text{ (M}^{-1}\text{s}^{-1}\text{)}$
Biotin-LVSR-AOMK	$9.1 (\pm 1.6) \times 10^3$
Cy5-LVSR-AOMK	$9.3 (\pm 1.6) \times 10^3$
Z-VRPR-FMK	$6.0 (\pm 0.7) \times 10^3$


SEs were derived from the linear regression analyses of the  $k_{\text{obs}}/I$  plots.

mammalian cell line transfected with the enzyme. MALT1 activity has been reported to be activated upon its co-overexpression with Bcl10 (Coornaert et al., 2008; Rebeaud et al., 2008). We therefore overexpressed MALT1 and Bcl10 in different combinations in human HEK293A cells, followed by cell lysis in the presence of probe, SDS-PAGE, and western blot analysis. As shown in Figure 3A, overexpression of MALT1-WT in combination with Bcl10 leads to Cy5 fluorescence of a protein species around 100 kDa. Overexpression of MALT1 alone shows fluorescence as well, albeit to a much lesser degree. Overexpression of Bcl10 alone or in the presence of MALT1-C464A likewise leads to the appearance of a faint fluorescently labeled protein. Note that the small size difference is due to the Flag-tag of the overexpression construct. No fluorescence can be seen when MALT1-C464A is expressed alone, again indicating that Cy5-LVSR-AOMK cannot bind the catalytic mutant but requires an intact active site. Furthermore, these data confirm that the probe has generally low cross-reactivity toward other proteins in the cell.

To inhibit binding of the probe to cathepsins, members of cysteine protease clan CA—some of which have substrate specificities that could overlap with MALT1—all samples contained the protease inhibitor E-64, which has been shown to potently inhibit cathepsins (Barrett et al., 1982). Under these experimental conditions, the probe bound most strongly to MALT1, and only a faint additional species around 31 kDa was seen. To determine whether this cross-reactivity would increase in the absence of E-64, we omitted the inhibitor from one sample. No significant intensity increase of the 31 kDa species could be seen, suggesting that the probe does not substantially cross-react with cathepsins, further validating its high degree of selectivity.

The biotin probe also enables the detection of active MALT1 after overexpression of MALT1-WT and Bcl10 (Figure 3B). No other probe-labeled proteins were detected, most likely due to the lower overall intensity of the biotin probe. Overall, it can be concluded that both probes can label and inhibit active MALT1 obtained from total cell lysates after overexpression in HEK293A cells.

The blot for MALT1 antigen reveals two species, one at approximately the same size as endogenous MALT1, which can be seen in the nontransfected samples, and one that migrates at approximately 10 kDa lower. The lower species is not detected by the Flag antibody if the Flag-tag is placed at the N instead of C terminus (Figure S4). In the WT overexpression samples, this protein species is labeled by the probe, suggesting that it represents an active C-terminal fragment. The fragment does not match the size of the species observed in the recombinant *E. coli*-produced samples (Figure 2). Again, both WT and mutant MALT1 are cleaved, and the question remains



**Figure 3. Cy5-LVSR-AOMK and Biotin-LVSR-AOMK Can Detect Active MALT1 after Overexpression**

(A and B) Different combinations of Bcl10, MALT1-WT, or C464A, respectively, were overexpressed in HEK293A cells. Cells were lysed in the presence of Cy5-LVSR-AOMK (A) or biotin-LVSR-AOMK (B), respectively, precleared, and subjected to SDS-PAGE, fluorescence scanning, and western blot analysis with the indicated antibodies. E-64 was added or omitted in the lysis buffer to control for cathepsin binding of the probe. Note that endogenous and Flag-tagged versions of the proteins migrate slightly differently during the SDS-PAGE. The asterisks mark nonspecific bands. Data are representative of three independent experiments.

whether the cleavage is autocatalytic or caused by another protease. It is theoretically possible that endogenous WT MALT1 performs the cleavage in the mutant samples. However, this seems unlikely because the cleavage occurs to the same extent in all samples. While we did not further investigate this phenomenon, we speculate that MALT1 is cleaved by another protease. Whether this is only the case when MALT1 is overexpressed or whether it also occurs at endogenous levels cannot be definitively concluded due to the low intensity of the endogenous protein.


### Cy5-LVSR-AOMK Can Detect Endogenous MALT1 Activity in OCI-Ly3 ABC-DLBCL Cells

Having established that MALT1 activity can be detected in an overexpression system, we wanted to determine whether the more sensitive probe Cy5-LVSR-AOMK can be used to detect MALT1 activity in ABC-DLBCL cell lines. These have previously been reported to exhibit constitutive MALT1 activity, whereas cell lines derived from the germinal center B cell-like DLBCL subtype (GCB-DLBCL) did not (Ferch et al., 2009; Hailfinger et al., 2009). As shown in Figure 4A, the probe was able to visualize MALT1 activity in the ABC-DLBCL cell line OCI-Ly3, but not the ABC-DLBCL cell line OCI-Ly10 nor the GCB-DLBCL cell lines SUDHL-4 and BJAB (note that the faint fluorescent signal visible around 97 kDa in lanes 2–4 does not match in size with the MALT1 species detected by the antibody). While we were surprised to see the difference between OCI-Ly3 and OCI-Ly10 cells, the result nonetheless proved that the probe was sensitive enough to detect endogenous MALT1 activity and directly confirmed the constitutive MALT1 activity of OCI-Ly3 cells shown indirectly in previous publications (Ferch et al., 2009; Hailfinger et al., 2009). We asked why we could not observe activity of MALT1 as claimed previously for OCI-Ly10 cells (Ferch et al.,

2009; Hailfinger et al., 2009). Along these lines, we observed a decrease in signal intensity generated by active MALT1 in the OCI-Ly3 cell line with increasing time in culture. Upon thawing of a fresh vial, the signal was back to its initial intensity (data not shown). This shows that culture conditions can lead to changes that affect MALT1 activity. While we could not solve the discrepancy between our results concerning MALT1 activity for the OCI-Ly10 cell line and previously published data, we speculate that culture conditions and subclones could well cause differences in MALT1 activity. Possibly, lymphoma-derived cells initially under the influence of MALT1 as a driver mutation become insensitive and acquire additional survival characteristics during prolonged passages. This speculation is consistent with our findings, but further clarification is beyond the scope of the current work. However, since our results show that the probe can detect active MALT1 if present at sufficient concentrations, such as in OCI-Ly3 cells or under overexpression conditions, it has the potential to help determine whether MALT1 is active in the cells and conditions investigated in a given study.

Several additional fluorescent bands (visible at lower molecular weights around 45, 31, and 14 kDa) were detected in all DLBCL lines, but their intensities were lower than that of the fluorescent MALT1 species at approximately 90 kDa. To investigate whether these species represent active cleavage products of MALT1 or nonspecific binding of the probe, we preincubated OCI-Ly3 cells with increasing concentrations of the reversible MALT1 inhibitor mepazine (Nagel et al., 2012; Schlauderer et al., 2013). This caused a dramatic decrease in the intensity of the fluorescent species at 90 kDa, while the intensity of the lower molecular weight species remained unchanged, suggesting that they represent nonspecific probe binding rather than MALT1 cleavage products. There is a formal possibility that the





**Figure 4. Cy5-LVSR-AOMK Can Detect Active Endogenous MALT1 in OCI-Ly3 Cells**

(A) The ABC-DLBCL cell lines OCI-Ly3 and OCI-Ly10 and the GCB-DLBCL cell lines SUDHL-4 and BJAB were lysed in the presence of Cy5-LVSR-AOMK and E-64, precleared, and subjected to SDS-PAGE, fluorescence scanning, and western blot analysis with the indicated antibodies. Note that the MALT1 band detected via Cy5 fluorescence in the OCI-Ly3 sample correlates with the band detected by the MALT1 antibody at ~90 kDa (lane 1). The faint fluorescent bands visible in lanes 2–4 do not correspond to the species detected by the antibody but migrate at a slightly higher molecular weight.

(B) OCI-Ly3 cells were treated with the indicated concentrations of mepazine for 4 hr, followed by lysis as in (A) except that the lysis buffer was supplemented with the appropriate concentrations of mepazine. Data are representative of three independent experiments. IB, immunoblot.


bands represent MALT1 cleavage fragments insensitive to mepazine. However, considering that the cross-reacting species are seen equally in all investigated DLBCL cell lines regardless of MALT1 activation status, it seems unlikely that the species are MALT1 cleavage products. The identity of the cross-reacting species was not investigated further at this point. Overall, Cy5-LVSR-AOMK was successfully used to detect active endogenous MALT1, and the constitutive activity of MALT1 in OCI-Ly3 ABC-DLBCL cells was confirmed in a direct manner.

#### Endogenous Active MALT1 Is Mostly NP40 Soluble, while Overexpression Leads to the Formation of NP40-Insoluble Aggregates

After establishing that the activity-based probes can label active MALT1 in cell lysates, we deployed the Cy5 probe to address questions of MALT1 biology. The localization and assembly of components of the CBM complex have been subjects of considerable research with varying results depending on activation status and overexpression versus endogenous proteins (Che et al., 2004; Gaide et al., 2002; Guiet and Vito, 2000; Izumiyama et al., 2003; Qiao et al., 2013; Rossman et al., 2006; Schaefer et al., 2004; Yan et al., 1999). Until now it has not been possible to differentiate between active material and total protein. To determine the subcellular location of MALT1 activation, we employed imaging flow cytometry and confocal microscopy. Our results were equivocal (data not shown), which we attribute to the hydrophobic nature of the Cy5 probe, such that even after extensive washing it remained partially bound to cells. This rendered it difficult to distinguish between cell penetration

and nonspecific binding to the outside of cells. We conclude that the Cy5 probe shows at best low cell penetration potential and that further optimization is needed if the probe is to be utilized for the detection and inhibition of MALT1 in intact cells.

Notwithstanding this limitation, we were able to obtain data on cell compartmentalization comparing endogenous versus overexpressed Bcl10 and MALT1 by employing differential detergent extraction. In the endogenous scenario, OCI-Ly3 cells constitutively activate MALT1 due to an activating mutation in CARMA1 (Lamason et al., 2010; Lenz et al., 2008). In the overexpression scenario, high protein concentrations of Bcl10 and MALT1 in HEK293A cells activate the protease, most likely by causing constitutive oligomerization. In OCI-Ly3 cells, the majority of endogenous MALT1, both active and total protein, as well as Bcl10, was detected in the NP40-soluble fraction (Figure 5A). This allowed us to conclude that the majority of active and total MALT1 is either present in soluble form in OCI-Ly3 cells or in a form that is solubilized under the lysis conditions used. In stark contrast to the OCI-Ly3 cell line, the overwhelming amount of active MALT1 was detected in the NP40-insoluble fraction upon ectopic coexpression of MALT1 and Bcl10 in HEK293A cells (Figure 5B). Bcl10 was divided relatively evenly between the NP40-soluble and -insoluble fractions, whereas MALT1 protein was only present at substantial amounts in the NP40-insoluble fraction when coexpressed with Bcl10. Importantly, when overexpressed alone, MALT1 remained soluble. Significant amounts of cleaved Bcl10 are seen only in the NP40-insoluble fraction of cells overexpressing both Bcl10 and MALT1-WT, confirming that probe binding correlates with



**Figure 5. Endogenous MALT1 Is NP40 Soluble, whereas Over-expressed Active MALT1 Is Mostly NP40 Insoluble**

(A and B) OCI-Ly3 cells (A) or HEK293A cells transfected as indicated (B) were lysed in the presence of Cy5-LVSR-AOMK, followed by centrifugation and separation of the lysates into NP40-soluble and -insoluble fractions, SDS-PAGE, fluorescence scanning, and western blot analysis with the indicated antibodies. Data are representative of three independent experiments.

MALT1 proteolytic activity. The fact that the majority of cleaved Bcl10 is detected in the NP40-insoluble fraction together with the vast majority of active MALT1 suggests that Bcl10 associated with active MALT1 is cleaved preferentially.

We conclude that upon overexpression Bcl10 forms NP40-insoluble aggregates that contain a large fraction of total Bcl10. MALT1, on the other hand, stays soluble when overexpressed alone, but is recruited to the aggregates upon co-overexpression with Bcl10. Whether active MALT1 seen in the NP40-soluble fraction originates from MALT1 that is active outside of the aggregates or whether the aggregates are partially solubilized during lysis releasing some of the active MALT1 could not be determined from the experimental setup. Because of the formation of NP40-insoluble aggregates, we conclude that total cell lysates should be used for analysis when studying the CBM complex to avoid missing crucial information.

### Mutation of MALT1 at K644 Decreases but Does Not Eliminate MALT1 Activity

As another example of probe application, we investigated the importance of ubiquitination for MALT1 activity. It has been reported that monoubiquitination of MALT1 at K644 is essential for its proteolytic activity (Pelzer et al., 2013). We created the ubiquitin mutant K644R and checked its effect on MALT1 activity using probe binding as a readout. While overexpression of the catalytic mutant C464A lowered Cy5 probe binding to endogenous levels, the K644R mutant still bound the probe. It was substantially decreased compared with MALT1-WT, but ubiquitination at K644 was clearly not a requirement for MALT1 activity per se, which was supported by the decreased but clearly detectable level of Bcl10 cleavage seen in the K644R sample.


### DISCUSSION

We have described probes that can distinguish between active and inactive MALT1 both in the recombinant in vitro setting and in cell lysates.

After purification of MALT1 from *E. coli*, two distinct species are visible upon SDS-PAGE analysis that correspond to full-length MALT1 and a C-terminal fragment, respectively (Figures 2 and S3). Both represent active species of MALT1 and show probe binding. It is interesting to note that the fluorescence intensity of the two protein species is approximately equal in the sodium citrate containing samples even though the upper species is more abundant, as shown by the coomassie stain. In the absence of citrate, labeling of the C-terminal fragment is strongly decreased compared with full-length enzyme. We speculate that this could be due to the absence of the N-terminal domains in this fragment, which have been reported to self oligomerize (Qiu and Dhe-Paganon, 2011). In the presence of sodium citrate, these domains might not play as important a role as in its absence.

Comparing the localization of MALT1 and Bcl10 protein in endogenous versus overexpression conditions revealed a clear discrepancy (Figure 5). While overexpressed active MALT1 localized predominantly in the NP40-insoluble fraction, active endogenous MALT1 was detected mostly in the NP40-soluble fraction. These results provide an example of how overexpression may create artificial conditions that do not accurately represent endogenous cellular processes. It remains to be determined whether accumulation in the NP40-insoluble fraction is simply due to higher protein levels upon overexpression, driving the formation of larger complexes that reach a critical size and thus become NP40 insoluble, or whether an entirely different mechanism of activation occurs under the different conditions. In case of the latter, it would be interesting to reexamine the recent results pertaining to the filamentous aggregate structure of the CBM complex (Qiao et al., 2013).

Considering the NP40-insoluble nature of overexpressed active MALT1, we reevaluated the results shown in Figure 3 for which NP40-soluble material only had been analyzed. Even though we had to conclude that the amount of active MALT1 was most likely underestimated due to the experimental conditions, the main conclusion—that both Cy5-LVSR-AOMK and biotin-LVSR-AOMK can detect active MALT1 and distinguish



**Figure 6. Mutation of MALT1 at K644 Decreases Its Activity**

HEK293A cells were transfected as indicated and lysed in the presence of Cy5-LVSR-AOMK, followed by SDS-PAGE of total cell lysates (no separation into NP40-soluble and -insoluble fractions), fluorescence scanning, and western blot analysis with the indicated antibodies. Note that the antibodies detect both endogenous and Flag-tagged versions of the proteins, which migrate at a slightly different molecular weight. In addition, the MALT1 antibody detects MALT1 cleavage products of unknown significance, and the Bcl10 antibody detects multiple higher molecular weight species, which most likely represent posttranslationally modified variants. Data are representative of three independent experiments.

between active and inactive MALT1 under overexpression conditions—remains valid.

When we elucidated the significance of MALT1 mutation at K644, which has been reported to alter a ubiquitination site important for its proteolytic activity (Pelzer et al., 2013), we determined that while the mutation clearly decreases MALT1 activity, ubiquitination at this site does not seem to be an absolute requirement (Figure 6). It is important to note that we could not detect a distinct higher molecular weight species representing ubiquitinated MALT1, which could be due to a lack of resolution during SDS-PAGE analysis. Another explanation could be the activity of deubiquitinating enzymes (DUBs) during lysis. The detection of ubiquitinated protein species is challenging due to the highly reversible nature of the modification. For efficient detection the cysteine alkylator N-ethylmaleimide (NEM) is commonly employed to inhibit DUBs, but in our case, this was not feasible because NEM also inhibits the cysteine protease MALT1 and probe binding. Therefore, we might be slightly underestimating the significance of ubiquitination for proteolytic activity. However, the important point is that in the absence of ubiquitination—whether achieved through the K644R mutation or DUBs—the probe still detects active MALT1. This, in combination with the fact that Bcl10 cleavage is decreased but not eliminated upon expression of the K644R mutant, points to a supporting role for ubiquitination at K644, although possibly not as essential as suggested in a previous report (Pelzer et al., 2013).

Another interesting observation when comparing the K644R mutant and WT enzyme is that MALT1 seems to be partially cleaved, resulting in a slightly smaller fluorescent species than full-length enzyme. Compared with the K644R mutant, the

cleavage is preferentially seen in the WT enzyme, and no cleavage is detected at all in the catalytic mutants, whether they retain their ubiquitin binding site or not. This raises the exciting possibility that the cleavage could potentially be auto-catalytic. The difference in size is very small and more readily detected via fluorescence than by immunodetection. Probe fluorescence is therefore a powerful tool that can be used to detect small but potentially important effects involving MALT1 activity.

## SIGNIFICANCE

We have developed two activity-based probes that specifically bind and inhibit active MALT1. The probes can differentiate effectively between active and inactive MALT1 both when using recombinant enzyme, cellular overexpression, and endogenous MALT1 in an ABC-DLBCL cell line. We have therefore provided tools that allow the monitoring of MALT1 activity. As examples of probe application, we compared the subcellular localization of active MALT1 under endogenous and overexpression conditions as well as the influence of ubiquitin modification of MALT1 at K644 on its activity. We propose that these tools will prove to be useful in further elucidating MALT1 and the importance of its scaffolding versus proteolytic function.

## EXPERIMENTAL PROCEDURES

### Probe Synthesis

See the Supplemental Information for a detailed description of the probe synthesis.

### Protein Expression

Recombinant full-length MALT1 was expressed and purified as previously described (Hachmann et al., 2012).

### Plasmids

Full-length MALT1 optimized for human expression (WT and catalytic mutant C464A, respectively), and Bcl10 were cloned into pcDNA3.1 containing a C-terminal Flag-tag. The K644R mutant was obtained using site-directed mutagenesis and cloned into the same vector as MALT1-WT and C464A.

### Incubation of Recombinant MALT1 with Cy5-LVSR-AOMK

Recombinant MALT1 and Cy5-LVSR-AOMK were incubated 30 min at 37°C in MALT1 assay buffer (50 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid, 100 mM NaCl, 10 mM dithiothreitol (DTT), 1 mM EDTA [pH 7.5]) with or without 0.9 M sodium citrate, followed by trichloroacetic acid (TCA) precipitation and SDS-PAGE using 4%–12% Bolt Bis-Tris gels (Life Technologies). Fluorescence was scanned at 700 nm using an Odyssey infrared scanner (LI-COR Biosciences), and total protein was visualized using coomassie-based InstantBlue stain (expedion).

### Inhibition Kinetics

Recombinant full-length MALT1 was incubated with increasing concentrations of inhibitor in MALT1 assay buffer with 0.9 M sodium citrate in the presence of 100 μM of the synthetic tetrapeptide substrate Ac-LRSR-AFC (SM Biochemicals). Kinetics of inhibition were evaluated by monitoring the release of the fluorophore over time using a Gemini Molecular Devices microplate spectrofluorometer. Progress curve data were analyzed by least-squares analysis using GraphPad Prism to determine the pseudo first-order rate constant  $k_{obs}$ . Derived second-order rate constants ( $k_{obs}/I$ ) were calculated taking into account the factor  $(1 + S/K_m)$ , where  $S$  is the substrate concentration and  $K_m$  determined as 37.4 μM (Hachmann et al., 2012).

### Cell Culture and Transfection

HEK293A cells were cultured in Dulbecco's modified essential medium supplemented with 10% fetal bovine serum (FBS), L-glutamine, and antibiotics. Transfections were performed when the cells had reached a density of ~50% using the NanoJuice transfection reagent (Novagen) according to the manufacturer's instructions. Cells were cultured for 24 hr after transfection. The DLBCL cell lines OCI-Ly3, OCI-Ly10, SUDHL-4, and BJAB were a generous gift from the laboratory of John C. Reed, Sanford-Burnham Medical Research Institute. OCI-Ly3 and OCI-Ly10 were cultured in Iscove's modified Dulbecco's medium supplemented with 20% human AB serum (Corning), L-glutamine,  $\beta$ -mercaptoethanol, and antibiotics. SUDHL-4 and BJAB were cultured in RPMI 1640 medium supplemented with 10% FBS, L-glutamine,  $\beta$ -mercaptoethanol, and antibiotics.

### Cell Lysis and Inhibitor Treatment

At 24 hr after transfection, cells were trypsinized and washed with PBS before lysis with a buffer containing 10 mM Tris (pH 7.4), 75 mM NaCl, 5 mM EDTA, 1% Nonidet P-40 (NP40), supplemented with 10 mM DTT and protease inhibitors (E-64, MG-132, 3,4-DCI (3,4-dichloroisocoumarin), and leupeptin, all at 10  $\mu$ M) unless noted otherwise. For the detection of endogenous MALT1 in DLBCL cells, 2  $\times$  10<sup>6</sup> OCI-Ly3 cells or 6  $\times$  10<sup>6</sup> OCI-Ly10, SUDHL-4, or BJAB cells, respectively, were washed with PBS and lysed as described above. Where applicable, Cy5-LVSR-AOMK or biotin-LVSR-AOMK, respectively, were added to the lysis buffer at 1  $\mu$ M. Cells were lysed 15–30 min at 37°C, followed by centrifugation and separation into NP40-soluble and -insoluble fractions. Alternatively, in Figure 6, total cell lysates were obtained by adding SDS sample buffer to the uncleared lysates. Total cell lysates and NP40-insoluble fractions were sonicated to ensure complete solubilization. The lysates were resolved via SDS-PAGE, followed by fluorescence scanning at 700 nm using an Odyssey infrared scanner (LI-COR Biosciences), transfer onto nitrocellulose membrane, and antibody detection. Mepazine hydrochloride was a generous gift from the laboratory of Daniel Krappmann, Helmholtz Zentrum München-German Research Center for Environmental Health, Institute of Molecular Toxicology and Pharmacology.

### Antibodies

Anti-MALT1 antibody was from Santa Cruz (B-12). Anti-Bcl10 was from Cell Signaling (4237). Anti-Flag was from Sigma-Aldrich (clone M2). Anti- $\beta$ -tubulin was from Abcam (ab6046). IRDye 800CW Streptavidin (LI-COR Biosciences) was used for biotin detection.

### SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures and four figures and can be found with this article online at <http://dx.doi.org/10.1016/j.chembiol.2014.11.011>.

### AUTHOR CONTRIBUTIONS

J.H. designed and performed the experiments. M.B. and M.D. designed, L.E.E.-M., and M.P. designed and synthesized, and L.E.S. synthesized the probes. J.H., M.B., and G.S.S. conceived the study and wrote the paper.

### ACKNOWLEDGMENTS

We thank Daniel Krappmann and John C. Reed for supplying reagents. We also thank Khatereh Motamedchaboki (Sanford-Burnham Medical Research Institute, Proteomics Facility), supported by NIH grant P30 CA030199, for help with mass spectrometry. This research was supported by NIH grants R01-GM09040 and R01 CA163743 to G.S.S. and R01 EB005011 and R01 HL11630703 to M.B. M.D. received support from the Foundation for Polish Science and a statutory activity subsidy from the Polish Ministry of Science and Higher Education for the Faculty of Chemistry at Wrocław University of Technology.

Received: July 22, 2014

Revised: November 13, 2014

Accepted: November 16, 2014

Published: December 31, 2014

### REFERENCES

- Akagi, T., Motegi, M., Tamura, A., Suzuki, R., Hosokawa, Y., Suzuki, H., Ota, H., Nakamura, S., Morishima, Y., Taniwaki, M., and Seto, M. (1999). A novel gene, MALT1 at 18q21, is involved in t(11;18) (q21;q21) found in low-grade B-cell lymphoma of mucosa-associated lymphoid tissue. *Oncogene* 18, 5785–5794.
- Barrett, A.J., Kembhavi, A.A., Brown, M.A., Kirschke, H., Knight, C.G., Tamai, M., and Hanada, K. (1982). L-trans-Epoxy succinyl-leucylamido(4-guanidino) butane (E-64) and its analogues as inhibitors of cysteine proteinases including cathepsins B, H and L. *Biochem. J.* 201, 189–198.
- Berger, A.B., Sexton, K.B., and Bogoy, M. (2006). Commonly used caspase inhibitors designed based on substrate specificity profiles lack selectivity. *Cell Res.* 16, 961–963.
- Cabalzar, K., Pelzer, C., Wolf, A., Lenz, G., Iwaszkiewicz, J., Zoete, V., Haifinger, S., and Thome, M. (2013). Monoubiquitination and activity of the paracaspase MALT1 requires glutamate 549 in the dimerization interface. *PLoS ONE* 8, e72051.
- Che, T., You, Y., Wang, D., Tanner, M.J., Dixit, V.M., and Lin, X. (2004). MALT1/paracaspase is a signaling component downstream of CARMA1 and mediates T cell receptor-induced NF- $\kappa$ B activation. *J. Biol. Chem.* 279, 15870–15876.
- Coornaert, B., Baens, M., Heynckx, K., Bekaert, T., Haegman, M., Staal, J., Sun, L., Chen, Z.J., Marynen, P., and Beyaert, R. (2008). T cell antigen receptor stimulation induces MALT1 paracaspase-mediated cleavage of the NF- $\kappa$ B inhibitor A20. *Nat. Immunol.* 9, 263–271.
- Dierlamm, J., Baens, M., Wlodarska, I., Stefanova-Ouzounova, M., Hernandez, J.M., Hossfeld, D.K., De Wolf-Peeters, C., Hagemeijer, A., Van den Berghe, H., and Marynen, P. (1999). The apoptosis inhibitor gene API2 and a novel 18q gene, MLT, are recurrently rearranged in the t(11;18)(q21;q21) associated with mucosa-associated lymphoid tissue lymphomas. *Blood* 93, 3601–3609.
- Edgington, L.E., Verdoes, M., Ortega, A., Withana, N.P., Lee, J., Syed, S., Bachmann, M.H., Blum, G., and Bogoy, M. (2013). Functional imaging of legumain in cancer using a new quenched activity-based probe. *J. Am. Chem. Soc.* 135, 174–182.
- Eitelhuber, A.C., Vasyka, O., Nagel, D., Bogner, M., Lenze, D., Lammens, K., Schlauderer, F., Hlahla, D., Hopfner, K.-P., Lenz, G., et al. (2014). Activity-based probes for detection of active MALT1 paracaspase in immune cells and lymphomas. *Chem Biol.* 22, this issue, 129–138.
- Ferch, U., Kloos, B., Gewies, A., Pfänder, V., Düwel, M., Peschel, C., Krappmann, D., and Ruland, J. (2009). Inhibition of MALT1 protease activity is selectively toxic for activated B cell-like diffuse large B cell lymphoma cells. *J. Exp. Med.* 206, 2313–2320.
- Gaide, O., Favier, B., Legler, D.F., Bonnet, D., Brissoni, B., Valitutti, S., Bron, C., Tschoep, J., and Thome, M. (2002). CARMA1 is a critical lipid raft-associated regulator of TCR-induced NF- $\kappa$ B activation. *Nat. Immunol.* 3, 836–843.
- Guert, C., and Vito, P. (2000). Caspase recruitment domain (CARD)-dependent cytoplasmic filaments mediate bcl10-induced NF- $\kappa$ B activation. *J. Cell Biol.* 148, 1131–1140.
- Hachmann, J., Snipas, S.J., van Raam, B.J., Cancino, E.M., Houlihan, E.J., Poreba, M., Kasperkiewicz, P., Drag, M., and Salvesen, G.S. (2012). Mechanism and specificity of the human paracaspase MALT1. *Biochem. J.* 443, 287–295.
- Haifinger, S., Lenz, G., Ngo, V., Posvitz-Fejfar, A., Rebeaud, F., Guzzardi, M., Penas, E.M., Dierlamm, J., Chan, W.C., Staudt, L.M., and Thome, M. (2009). Essential role of MALT1 protease activity in activated B cell-like diffuse large B-cell lymphoma. *Proc. Natl. Acad. Sci. USA* 106, 19946–19951.
- Haifinger, S., Nogai, H., Pelzer, C., Jaworski, M., Cabalzar, K., Charton, J.E., Guzzardi, M., Décaillat, C., Grau, M., Dörken, B., et al. (2011). Malt1-dependent RelB cleavage promotes canonical NF- $\kappa$ B activation in lymphocytes and lymphoma cell lines. *Proc. Natl. Acad. Sci. USA* 108, 14596–14601.
- Isaacson, P.G., and Du, M.Q. (2004). MALT lymphoma: from morphology to molecules. *Nat. Rev. Cancer* 4, 644–653.



- Izumiyama, K., Nakagawa, M., Yonezumi, M., Kasugai, Y., Suzuki, R., Suzuki, H., Tsuzuki, S., Hosokawa, Y., Asaka, M., and Seto, M. (2003). Stability and subcellular localization of API2-MALT1 chimeric protein involved in t(11;18)(q21;q21) MALT lymphoma. *Oncogene* 22, 8085–8092.
- Jeltsch, K.M., Hu, D., Brenner, S., Zöller, J., Heinz, G.A., Nagel, D., Vogel, K.U., Rehage, N., Warth, S.C., Edelmann, S.L., et al. (2014). Cleavage of roquin and regnase-1 by the paracaspase MALT1 releases their cooperatively repressed targets to promote TH17 differentiation. *Nat. Immunol.* 15, 1079–1089.
- Kato, D., Boatright, K.M., Berger, A.B., Nazif, T., Blum, G., Ryan, C., Chehade, K.A., Salvesen, G.S., and Bogoy, M. (2005a). Activity-based probes that target diverse cysteine protease families. *Nat. Chem. Biol.* 1, 33–38.
- Kato, D., Verhelst, S.H., Sexton, K.B., and Bogoy, M. (2005b). A general solid phase method for the preparation of diverse azapeptide probes directed against cysteine proteases. *Org. Lett.* 7, 5649–5652.
- Lamason, R.L., McCully, R.R., Lew, S.M., and Pomerantz, J.L. (2010). Oncogenic CARD11 mutations induce hyperactive signaling by disrupting autoinhibition by the PKC-responsive inhibitory domain. *Biochemistry* 49, 8240–8250.
- Lenz, G., Davis, R.E., Ngo, V.N., Lam, L., George, T.C., Wright, G.W., Dave, S.S., Zhao, H., Xu, W., Rosenwald, A., et al. (2008). Oncogenic CARD11 mutations in human diffuse large B cell lymphoma. *Science* 319, 1676–1679.
- Lucas, P.C., Yonezumi, M., Inohara, N., McAllister-Lucas, L.M., Abazeed, M.E., Chen, F.F., Yamaoka, S., Seto, M., and Nunez, G. (2001). Bcl10 and MALT1, independent targets of chromosomal translocation in malt lymphoma, cooperate in a novel NF-kappa B signaling pathway. *J. Biol. Chem.* 276, 19012–19019.
- Morgan, J.A., Yin, Y., Borowsky, A.D., Kuo, F., Nourmand, N., Koontz, J.I., Reynolds, C., Soreng, L., Griffin, C.A., Graeme-Cook, F., et al. (1999). Breakpoints of the t(11;18)(q21;q21) in mucosa-associated lymphoid tissue (MALT) lymphoma lie within or near the previously undescribed gene MALT1 in chromosome 18. *Cancer Res.* 59, 6205–6213.
- Nagel, D., Spranger, S., Vincendeau, M., Grau, M., Raffegerst, S., Kloos, B., Hlahla, D., Neuenschwander, M., Peter von Kries, J., Hadian, K., et al. (2012). Pharmacologic inhibition of MALT1 protease by phenothiazines as a therapeutic approach for the treatment of aggressive ABC-DLBCL. *Cancer Cell* 22, 825–837.
- Oeckinghaus, A., Wegener, E., Welteke, V., Ferch, U., Arslan, S.C., Ruland, J., Scheidereit, C., and Krappmann, D. (2007). Malt1 ubiquitination triggers NF-kappaB signaling upon T-cell activation. *EMBO J.* 26, 4634–4645.
- Pelzer, C., Cabalzar, K., Wolf, A., Gonzalez, M., Lenz, G., and Thome, M. (2013). The protease activity of the paracaspase MALT1 is controlled by mono-ubiquitination. *Nat. Immunol.* 14, 337–345.
- Pomerantz, J.L., Denny, E.M., and Baltimore, D. (2002). CARD11 mediates factor-specific activation of NF-kappaB by the T cell receptor complex. *EMBO J.* 21, 5184–5194.
- Powers, J.C., Asgjan, J.L., Ekici, O.D., and James, K.E. (2002). Irreversible inhibitors of serine, cysteine, and threonine proteases. *Chem. Rev.* 102, 4639–4750.
- Qiao, Q., Yang, C., Zheng, C., Fontán, L., David, L., Yu, X., Bracken, C., Rosen, M., Melnick, A., Egelman, E.H., and Wu, H. (2013). Structural architecture of the CARMA1/Bcl10/MALT1 signalosome: nucleation-induced filamentous assembly. *Mol. Cell* 51, 766–779.
- Qiu, L., and Dhe-Paganon, S. (2011). Oligomeric structure of the MALT1 tandem Ig-like domains. *PLoS ONE* 6, e23220.
- Rebeaud, F., Haifinger, S., Posevitz-Fejfar, A., Tapernoux, M., Moser, R., Rueda, D., Gaide, O., Guzzardi, M., Iancu, E.M., Rufer, N., et al. (2008). The proteolytic activity of the paracaspase MALT1 is key in T cell activation. *Nat. Immunol.* 9, 272–281.
- Rosebeck, S., Madden, L., Jin, X., Gu, S., Apel, I.J., Appert, A., Hamoudi, R.A., Noels, H., Sagaert, X., Van Loo, P., et al. (2011). Cleavage of NIK by the API2-MALT1 fusion oncoprotein leads to noncanonical NF-kappaB activation. *Science* 331, 468–472.
- Rossman, J.S., Stoicheva, N.G., Langel, F.D., Patterson, G.H., Lippincott-Schwartz, J., and Schaefer, B.C. (2006). POLKADOTS are foci of functional interactions in T-Cell receptor-mediated signaling to NF-kappaB. *Mol. Biol. Cell* 17, 2166–2176.
- Ruefli-Brasse, A.A., French, D.M., and Dixit, V.M. (2003). Regulation of NF-kappaB-dependent lymphocyte activation and development by paracaspase. *Science* 302, 1581–1584.
- Ruland, J., Duncan, G.S., Wakeham, A., and Mak, T.W. (2003). Differential requirement for Malt1 in T and B cell antigen receptor signaling. *Immunity* 19, 749–758.
- Schaefer, B.C., Kappler, J.W., Kupfer, A., and Marrack, P. (2004). Complex and dynamic redistribution of NF-kappaB signaling intermediates in response to T cell receptor stimulation. *Proc. Natl. Acad. Sci. USA* 101, 1004–1009.
- Schlauderer, F., Lammens, K., Nagel, D., Vincendeau, M., Eitelhuber, A.C., Verhelst, S.H., Kling, D., Chrusciel, A., Ruland, J., Krappmann, D., and Hopfner, K.P. (2013). Structural analysis of phenothiazine derivatives as allosteric inhibitors of the MALT1 paracaspase. *Angew. Chem. Int. Ed. Engl.* 52, 10384–10387.
- Staal, J., Driege, Y., Bekaert, T., Demeyer, A., Muyliaert, D., Van Damme, P., Gevaert, K., and Beyaert, R. (2011). T-cell receptor-induced JNK activation requires proteolytic inactivation of CYLD by MALT1. *EMBO J.* 30, 1742–1752.
- Sun, L., Deng, L., Ea, C.K., Xia, Z.P., and Chen, Z.J. (2004). The TRAF6 ubiquitin ligase and TAK1 kinase mediate IKK activation by BCL10 and MALT1 in T lymphocytes. *Mol. Cell* 14, 289–301.
- Uehata, T., Iwasaki, H., Vandenbon, A., Matsushita, K., Hernandez-Cuellar, E., Kuniyoshi, K., Satoh, T., Mino, T., Suzuki, Y., Standley, D.M., et al. (2013). Malt1-induced cleavage of regnase-1 in CD4(+) helper T cells regulates immune activation. *Cell* 153, 1036–1049.
- Uren, A.G., O'Rourke, K., Aravind, L.A., Pisabarro, M.T., Seshagiri, S., Koonin, E.V., and Dixit, V.M. (2000). Identification of paracaspases and metacaspases: two ancient families of caspase-like proteins, one of which plays a key role in MALT lymphoma. *Mol. Cell* 6, 961–967.
- Wang, D., You, Y., Case, S.M., McAllister-Lucas, L.M., Wang, L., DiStefano, P.S., Nuñez, G., Bertin, J., and Lin, X. (2002). A requirement for CARMA1 in TCR-induced NF-kappa B activation. *Nat. Immunol.* 3, 830–835.
- Wegener, E., and Krappmann, D. (2007). CARD-Bcl10-Malt1 signalosomes: missing link to NF-kappaB. *Sci. STKE* 2007, pe21.
- Wiesmann, C., Leder, L., Blank, J., Bernardi, A., Melkko, S., Decock, A., D'Arcy, A., Villard, F., Erbel, P., Hughes, N., et al. (2012). Structural determinants of MALT1 protease activity. *J. Mol. Biol.* 419, 4–21.
- Yan, M., Lee, J., Schilbach, S., Goddard, A., and Dixit, V. (1999). mE10, a novel caspase recruitment domain-containing proapoptotic molecule. *J. Biol. Chem.* 274, 10287–10292.
- Yu, J.W., Jeffrey, P.D., Ha, J.Y., Yang, X., and Shi, Y. (2011). Crystal structure of the mucosa-associated lymphoid tissue lymphoma translocation 1 (MALT1) paracaspase region. *Proc. Natl. Acad. Sci. USA* 108, 21004–21009.
- Zhou, H., Wertz, I., O'Rourke, K., Ultsch, M., Seshagiri, S., Eby, M., Xiao, W., and Dixit, V.M. (2004). Bcl10 activates the NF-kappaB pathway through ubiquitination of NEMO. *Nature* 427, 167–171.