

Acute Pain Management

Rita Agarwal MD
Clinical Professor of Anesthesiology
Stanford University School of Medicine

Introduction

- The need for pain management in children is well accepted
- Children have frequently been under treated
- Education and research have lead to improved pain management
- Slides:
<http://med.stanford.edu/content/dam/sm/pedsanesthesia/documents/acute-pain-managment-2019.pdf>




Assessment

- Pain is difficult to assess due to:
 - Age, developmental level, maturity
 - Fear, anxiety, depression
 - Behavioral problems, learning disabilities
- Multiple available pain scales




Wong Baker Scales



Most commonly used at LPCH,

OUCHER SCALES



https://www.researchgate.net/publication/320001571_Sedation_and_analgesia_for_procedures_in_the_pediatriac_emergency_room/figures?lo=1

Faces Legs Activity Cry Consolability

FLACC Scale ²		0	1	2
1	Face	No particular expression or smile.	Occasional grimace or frown, withdrawn, disinterested.	Frequent to constant frown, clenched jaw, quivering chin.
2	Legs	Normal position or relaxed.	Uneasy, restless, tense.	Kicking, or legs drawn up.
3	Activity	Lying quietly, normal position, moves easily.	Squirming, shifting back and forth, tense.	Arched, rigid or jerking.
4	Cry	No crying (awake or asleep).	Moans or whimpers; occasional complaint.	Crying steadily, screams or sobs, frequent complaints.
5	Consolability	Content, relaxed.	Reassured by occasional touching, hugging or being talked to, distractible.	Difficult to console or comfort.

REFERENCES:

1. Pain FACES based on Wong D.L, Hockenberry-Eaton M, Wilson D, Winkelstein M.L, Schwatz P: *Wong's Essentials of Pediatric Nursing*, ed 6, St. Louis, 2001, p. 1301 © by Mosby, Inc.
2. From The FLACC: A behavioral scale for scoring postoperative pain in young children, by S Merkel and others, 1997, *Pediatr Nurse* 23(3), p. 293-297. ©1997 by Jannetti Co. University of Michigan Medical Center.
3. All other content and design ©Allen Perri Design Group, Ltd. DBA Healthcare Inspirations. All rights reserved.

Product ID: PGPA-130
(877) 646-5877
HealthcareInspirations.com/pain

Healthcare Inspirations 

Developed for use in patients post op, who are non-verbal, has been validated , in a variety of different pain populations.

Management Principles

Preoperative

- Prepare child and family for surgery
- Discuss past pain experiences (child and parent's)
- Present pain treatment options
- Discuss pain assessment tools

Management Principles

Postoperative

- Maximize efficacy of technique(s)
 - Appropriate & frequent pain assessment
 - Allow parents to help assess preverbal children
 - Combine techniques

Introduction




- Regional anesthesia in children is practical, safe and effective
- Children tend to have easier anatomy

Introduction

- Most regional techniques are placed in an anesthetized patient
- Always consider risk/benefit
- Calculate toxic dose
- “Perfect” block is not mandatory
- Optimize use of other medications




Neuraxial Blocks



Caudal Analgesia


- One shot or continuous catheters
- The sacrum is cartilaginous in infants and children, ossification occurs between the ages of 20 and 30 years.
- Malformations of the sacrum occur in ~ 10% of the population.
- The sacral cornua marks lateral edges of the sacral hiatus
- Always above the gluteal crease

Caudal Anatomy




Caudal Analgesia

- Insert needle in sacral hiatus at $\sim 60^\circ$ angle until LOR as sacrococcygeal membrane is pierced
- Drop needle until it is parallel to back and advance a few mm
- Aspirate and inject LA



Complications

- Intraosseous, intravenous or subcutaneous injection
- Coccygeal cornua



Suggested single Shot Dosing for Caudals*

Armitage	ml/kg	Approx level
	0.5	sacrolumbar
	0.75	Lumbar-mid thoracic
	1	mid –high thoracic
	1.25	high thoracic
Takasaki	0.06ml/seg/kg	

Toxic dose for B and LB: 1.5 mg/kg for neonates

2.5 mg/kg for children

2-3 mg/kg for R

*Rarely need to administer more than 15-20 mls

Test Dosing

- Although not 100% accurate
- Epinephrine 0.5 ug/kg should be used
- ST-T wave Δ 's, HR \uparrow by 10bpm, or increased BP are predicative of intravascular injections
- Isoproterenol and lower doses of epi do not seem to be effective
- ? If U/S decreases intravascular injection


Table 1. Summary of Studies of a Simulated Test Dose (IV Administration) in Pediatric-Aged Patients

Reference	Anesthetic	Test drug ($\mu\text{g}/\text{kg}$)	Criteria evaluated	Findings
Desparmet et al. (7)	1 MAC halothane 50% nitrous oxide/oxygen Atropine—randomized	Epinephrine 0.5 in lidocaine 1 mg/kg	HR, % change in SBP	Positive HR response: ≥ 10 bpm. Atropine increases HR sensitivity.
Perillo et al. (11)	1.2 MAC halothane 50% nitrous oxide/oxygen No atropine	Isoproterenol 0.05, 0.075 in 0.5 mg/kg lidocaine	HR ≥ 10 bpm	Higher dose more sensitive; 17/21 with 0.05 $\mu\text{g}/\text{kg}$ versus 22/23 with 0.075 $\mu\text{g}/\text{kg}$ I.
Kozek-Langenecker et al. (12)	Anesthetized & awake state; 1.2 MAC halothane 70% nitrous oxide/oxygen No atropine	Isoproterenol 0.05, 0.075 or 0.1 in 0.25 mg/kg bupivacaine	HR ≥ 20 bpm	All 3 doses of I. effective in awake state. Higher dose more sensitive during anesthesia.
Tanaka and Nishikawa (14)	1 MAC sevoflurane 60% nitrous oxide/oxygen Atropine—randomized	Epinephrine 0.5 in 1 mg/kg lidocaine	HR ≥ 10 bpm SBP ≥ 15 mm Hg	HR: 100% sensitive (15/15) with and without atropine. SBP—100% sensitive with atropine and 10/15 without atropine.
Sethna et al. (15)	1 MAC isoflurane 100% oxygen Atropine	Epinephrine 0.5, 0.75 in 1 mg/kg lidocaine	HR ≥ 10 bpm SBP ≥ 15 mm Hg	HR: 19/21 with 0.5 $\mu\text{g}/\text{kg}$ and 21/21 with 0.75 $\mu\text{g}/\text{kg}$. SBP: 17/21 with 0.5 $\mu\text{g}/\text{kg}$ and 19/21 with 0.75 $\mu\text{g}/\text{kg}$.
Kozek-Langenecker et al. (16)	1 MAC sevoflurane or halothane 70% nitrous oxide/oxygen No atropine	Incremental doses of isoproterenol in saline	HR ≥ 20 bpm	Higher dose of isoproterenol (55 vs 32 ng/kg) needed with sevoflurane than with halothane.
Tanaka and Nishikawa (22)	1 MAC sevoflurane 67% nitrous oxide/oxygen Atropine	Epinephrine 0.5 in 1 mg/kg lidocaine	HR ≥ 10 bpm SBP ≥ 15 mm Hg T wave $\geq 25\%$	Positive response in 16/16, 13/16, 16/16 with HR, SBP, T-wave respectively. Increase in T-wave amplitude occurred earliest.
Kozek-Langenecker et al. (23)	1 MAC sevoflurane or halothane 70% nitrous oxide/oxygen No atropine	Epinephrine 0.5 in saline	HR ≥ 10 bpm SBP ≥ 15 mm Hg T wave $\geq 25\%$	T-wave more sensitive than HR or SBP. T wave, SBP more sensitive with sevoflurane than with halothane.
Tanaka et al. (25)	1 MAC sevoflurane 67% nitrous oxide/oxygen Atropine	Epinephrine 0.5 in bupivacaine 0.25 mg/kg or I. In lidocaine 1 mg/kg	HR ≥ 10 bpm T wave $\geq 25\%$	HR and T-wave 100% sensitive with epinephrine. HR 100% sensitive with I. No change in T-wave noted with I.
Tanaka and Nishikawa (26)	1 MAC sevoflurane 67% nitrous oxide/oxygen Atropine	Epinephrine 0.125, 0.25, 0.5 in 0.25, 0.5, and 1 mg/kg lidocaine	HR ≥ 10 bpm SBP ≥ 15 mm Hg T wave $\geq 25\%$	0.125 $\mu\text{g}/\text{kg}$ not sensitive. 0.25, 0.5—100% sensitive for T-wave criteria. 0.5— 100% for HR criteria.

Lumbar and Thoracic Epidurals

• ANATOMY



- Dural sac extends to S₄ in the neonate
- Spinal cord ends at L₃₋₄
- Adult levels of S₃ & L₁ are reached by age 1 yr.
- Epidural contents are less dense
- Local anesthesia can spread easily



Lumbar and Thoracic Epidurals

• TECHNIQUE AND EQUIPMENT

- Same as adults
- Shorter, smaller needles and catheters are available
- Bromage grip
- Usually anesthetized child



Lumbar and Thoracic Epidurals

- DISTANCE TO EPIDURAL SPACE

- 0.5 - 1 cm neonates and infants

- 1 - 2 cm 1 - 5 yr.

- 2 - 3 cm 5 - 10 yr.

- 3 - 5 cm 10 - 16 yr.

Distance (mm) = (age (yrs) x 2) + 10

Distance (mm) = (weight x 0.8)

Lumbar and Thoracic Epidurals

- DOSING

- Toxic Dose of Bupivacaine
 - 0.5 mg/kg/hr in children
 - 0.2- 0.25 mg/kg/hr in neonates
- Ropivacaine is similar-
 - 0.6mg/kg/hr in children
 - 0.3 mg/kg/hr in neoantes
- This represents maximum dosing, NOT necessarily starting doses
- Children often do better with more volume

Lumbar and Thoracic Epidurals: Recommended Maximum or toxic dose

Agents	Single Shot Bolus
Bupivacaine	2-2.5 mg/kg
Ropivacaine	3 mg/kg
Fentanyl	1-2 ug/kg
Morphine	20-50 ug/kg
Hydromorphone	10 ug/kg

I usually start with:

0.2-0.3ml/kg for a lumbar epidural bolus dosing, based on IBW and
0.15-0.2 ml/kg for thoracic epidural dosing

Recommended starting doses for continuous or intermittent bolus dosing for epidural catheters

Agents	Conc	Infusion or Intermittent Epidural Bolus
Bupivacaine	0.08-0.125%	0.15-0.4 cc/kg/hr
Ropivacaine	0.08-0.2%	0.15-0.3 cc/kg/hr
Fentanyl	2-5 ug/cc	0.5-1 ug/kg/hr
Morphine	50-100 ug/cc	2-10 ug/kg/hr
Hydromorph	3-5ug/cc	1-3 ug/kg/hr

What's in a Stereo-isomer?

- Ropivacaine (R) and levobupivacaine (LB) were introduced to ↓ cardiotoxicity
- R is less potent and the least toxic, it may have less motor block
- LB seems to have equipotent to B but is less toxic (it is not available in the USA)
- Ease of resuscitation after overdose: $R > LB > B$
- R and LB < arrhythmogenic than B

A Pinch Of This

- Fentanyl
 - Safe, effective and popular in combination with local anesthetic in continuous epidural infusions
 - Most studies find no benefit to adding fentanyl to a one shot caudal for post-operative analgesia
 - Increased side effects

A Pinch of This.....

- Clonidine
 - Most studies show a prolongation of analgesia and increased sedation
 - Hypotension and bradycardia do not seem to occur as much in children
 - De Mey found no effect from either clonidine or sufentanil
 - Single shot 1-2 ug/kg
 - Continuous/IEB 0.1ug-1ug/cc solution. Most of us at LPCH use 0.1 or 0.2 ug/ml

De Mey JC et.al Eur J
Anaesth 17: 379-82, 2000

A Pinch of This.....

- De Negri and colleagues found the addition of ketamine 0.5mg/kg to be as effective as 2ug/kg clonidine when added to a 1 shot caudal
- Marhofer found ketamine 1mg/kg to provide equivalent analgesia to 0.25% bupivacaine

DeNegri et.al Paed Anaesth 11: 679-83,2001

Marhofer P et.al. Br J Anaesth 84: 341-5, 2000



Not approved in
US


Threading catheters

- In infants catheters can be threaded from the caudal space
- Coiling is more common with increasing age
- The usual complications can still occur
- Position should always be confirmed epidurogram is the gold standard , but increasingly U/S is being used
- Houck found 32% malposition
 - 10/28--too high
 - 17/28 coiled in lumbosacral area
 - 1 outside of epidural space

Epidurograms




Too Low




Just Right

• Too High




Just right



Opioids

- The mainstay of pain medications
- A variety of routes are available—IM is avoided
- Intravenous opioids are used in doses similar to adults
- Intermittent administration is easy and allows for frequent dose and/or interval changes
- Meperidine is associated with \uparrow neurologic symptoms



Opioid Analgesia

Constant Infusion

- Neonates, infants, developmentally delayed children
- Avoids peaks and troughs
- Useful for post-operative pain
- Morphine, hydromorphone or fentanyl are most commonly used

Opioid Analgesia

Patient Controlled Analgesia (PCA)

- Pts ~ 5 yr. and developmentally appropriate
- Parent and nurse controlled analgesia may be appropriate in selected cases
- With or without basal infusion ?
- Hydromorphone, morphine or fentanyl most commonly used
- May use Parent/Provider Controlled analgesia in selected patients



Patient Controlled Analgesia

Basal infusion

- Better pain control, improved sleep patterns for first 24 hr.
- Increased total narcotic usage
- increased reported side effects



Continuous Basal Infusion- Pro

- *Berde CB J Peds 1991; 118:460*
 - Orthopedic surgery
 - 3 groups: IM morphine, PCA bolus only, PCA bolus + CBI - Similar morphine use, side effects in all groups
 - Lowest pain scores in CBI group
- *Skues MA Ped Anesth 1993; 3:223*
 - Abdominal surgery
 - Similar morphine use in Bolus only and Bolus +CBI - Improved sleep in CBI group
- *Yildiz K. Ped Anesth 2003; 13:427*
 - Appendectomy
 - Higher Demerol consumption in PCA bolus only group
 - Trend toward improved pain scores in Bolus + CBI group - Similar side effects in both groups
- *Doyle E. BJA 1993; 71:818*
 - Appendectomy
 - 3 groups: Bolus only, Bolus + 4 mcg/kg/hr, Bolus + 10 mcg/kg/hr - Similar pain scores in all groups
 - Improved sleep in both CBI groups
 - 4 mcg/kg/hr : least number of hypoxemic episodes (SpO2 < 94%) - 10 mcg/kg/hr : greatest morphine use, PONV and hypoxemia

Continuous Basal Infusion- Com

- *Doyle E. BJA 1993; 71: 670*
- - Appendectomy
- - Bolus vs. Bolus + 20 mcg/kg/hr CBI
- - CBI group – Increased morphine use, sedation, PONV, hypoxemic episodes, better sleep
- *McNeely J. J Pain Symptom Manage 1997; 13: 268*
- - Lower extremity surgery
- - Bolus vs. Bolus + Nighttime CBI
- - Increased morphine use and hypoxemic episodes (SpO₂ < 90%) in CBI group

PCA- Opioid Dosing

Drug	Bolus Dose ($\mu\text{g}/\text{kg}$)	Lockout Interval (min)	CBI ($\mu\text{g}/\text{kg}/\text{hour}$)	4-hr. limit ($\mu\text{g}/\text{kg}$)
Morphine	10-20	8-15	0-20	250-400
Hydromorphone	2-4	8-15	0-4	50-80
Fentanyl	0.5	5-10	0-0.5	7-10

Initial dosing recommendations in opioid-naïve children

*Reproduced from Malviya S, Polaner DM, Berde CB. Acute Pain in Cote CJ, Lerman J, Anderson B eds
A Practice of Anesthesia for Infants and Children. Elsevier Inc 2013 pp 928-933*

Opioids

- Intranasal (IN) fentanyl
2ug/kg improves emergence
in children undergoing T&T
- ED now using IN sufentanil
?dose (0.7-2 ug/kg)
- IN butorphanol 25ug/kg
- IN remifentanil 2-4 ug/kg for
intubation



Oral Opioids

- Codeine 1mg/kg q 3-4 hrs FDA warning
- Hydrocodone 0.1-0.2 mg/kg q 3-4 hrs FDA warning
- Oxycodone 0.1-0.15 mg/kg q 3-4 hrs
- Most preparations come in an elixir as well as tablets, +/- acetaminophen

FDA Black
Box Warning
for Codeine
and Tramadol



Non-Opioid Analgesics

- Ibuprofen-5-10mg/kg po every 6-8 hours
- Ketorolac 0.5mg/kg IV every 8 hours up to 30mg
- Acetaminophen
 - Rectal 40mg/kg then 20mg/kg every 6 hrs
 - Oral 10-15mg/kg every 6 hours
 - IV 10-15 mg q 6 hours

How much opioid?

- 87% of pediatric patients received a prescription
- 60% used.
- Most families had NOT been counselled on safe disposal.
- Only 5% of families disposed of unused medications properly,
- 50% of families had teenagers in the house.
- After two weeks, families had on average 36 tablets (range 0-95) or 67 ml (range 0-567 ml) of prescription opioids left over.

How much opioid?

- Another study in children receiving morphine after surgery in Montreal noted that 1431 doses were ordered, and only 131 doses (9.2%) were administered
- Most were counselled, ~60% returned unused medication, ~25% disposed properly
- However, majority kept the medications out in the open

Opioid Prescribing for the Treatment of Acute Pain in Children on Hospital Discharge.

Figure 1





Figure 1 . Doses dispensed and consumed by surgical specialty..

*General pediatric surgery excluding Nuss procedure. **Orthopedic surgery excluding posterior spinal fusion. Card indicates cardiothoracic surgery; GU, genitourologic surgery; Neuro, neurosurgery; Pectus, Nuss procedure; Plastic, plastic surgery; PSF, posterior spinal fusion.




Almost **18 out of 100** Americans have used illicit drugs or misused Rx drugs.

Overdose Deaths Involving Opioids, by Type of Opioid, United States, 2000-2016



SOURCE: CDC/NCHS, National Vital Statistics System, Mortality; CDC WONDER, Atlanta, GA: US Department of Health and Human Services, CDC, 2017. <https://wonder.cdc.gov/>

Figure 2



Opioid Prescribing for the Treatment of Acute Pain in Children on Hospital Discharge.

Monitto, Constance et. al Anesthesia & Analgesia

Figure 2 . Duration of opioid use by specialty. Duration of opioid therapy reported by families after hospital discharge. Data are stratified by primary service.

Opioid Prescribing A&A Dec 2017

Calculating the cumulative amount of leftover opioid among

- 235 respondents:
 - 3110 oxycodone tabs
 - 7264 ml oxy elixir
 - Morphine, hydromorphone etc
 - 45,573 morphine mg equivalents
 - 19% advised on disposal
 - 4% disposed leftover opioids



Opioid Prescribing A&A 2017

- Girls use 7.5 X more opioids
- Ortho and Nuss bar highest opioid use post surgery
- Patient with pain scores >5




Who gets addicted?

Risk Factors for Prescription Opioid Pain Reliever Abuse and Overdose



Obtaining overlapping prescriptions from multiple providers and pharmacies.



Taking high daily dosages of prescription opioid pain relievers.



Having mental illness or a history of alcohol or other substance abuse.




Living in rural areas and having low income.

- Anyone can get addicted
- Post surgery 0.9 -12% will become chronic opioid users
- H/o psychiatric disease, including ADHD, depression, anxiety
- Lives in rural area
- Low income

What can we do?

- Educate EVERYONE-physicians, patients, dentists, nurses, physician assistants
- Research-how much pain? For how long? What else can be used?
- Mental Health resources
- Recognize early-treat quickly
- Limit supply—but what about street drugs?
- Prescription Drug Monitoring Programs



Created posters for
Practitioners

AAP News

October 20, 2017

Poster created to promote safe storage, disposal of Rx

Lucien Gonzalez, M.D., M.S., FAAP, Rita Agarwal, M.D., FAAP and Constance S. Houck, M.D., M.P.H., FAAP

Safe storage and disposal of controlled prescription drugs may save a young person's life.
It's that simple.

Medicine Safety for Children and Teens: We All Play a Role

DO NOT share prescribed medicines with anyone, including family members.

DO NOT save prescribed medicines, unless told to do so by your doctor.

DO secure all medicines up and out of reach of children and teens.

DO make sure children and teens take their medicines correctly.

DO follow all the instructions from your doctor or pharmacist.

DO talk with your child's doctor if you have any questions.

DO get rid of all old or unused medicines.

- Follow the instructions on the medicine label or package insert. Only flush medicines if the label says it is okay to do so.
- If the label doesn't give instructions, look for a "take back program" in your community.
- If instructions and "take back programs" are not available, take the medicine out of the original container and mix it with used coffee grounds, dirt, or kitty litter, and throw in the trash.
- Visit www.healthychildren.org/medicinesafety for details



American Academy
of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN



healthychildren.org
POWERED BY SPECIALISTS. TRUSTED BY PARENTS.
FROM THE AMERICAN ACADEMY OF PEDIATRICS



PCSS

Providers
Clinical Support
System

Funding for this initiative was made possible in part by grant no. 5R15P0200040-01 from SAMHSA. The views expressed in written or printed educational materials do not necessarily reflect the official policies of the Department of Health and Human Services, nor does mention of trade names, commercial practices, or organizations imply endorsement by the U.S. Government.

PCSS-O



Please Note: CME is only available for live events. CME is not available for recorded webinars.

Providers' Clinical Support System—Opioid Therapies (PCSS-O)

Start Date: September 2014

End Date: September 2017

Welcome, you are signed in as Rita Agarwal [Sign Out](#)

[Pediatrician View](#) [Program View](#) [Resident View](#)

[Home](#) [Continuing Education](#) [Learning Plan](#) [PediaLink QI](#) [Maintenance of Certification](#) [Learning Resources](#) [Leadership](#)

[My CME](#) [CME Transcript](#) [CME Finder](#) [About AAP CME](#) [Print](#)

My CME

A list of CME activities you are currently registered for through the AAP.

CME Activity	Start	End date	Activity Format	Claim credit expiration date	Action
Registered CME					
2017 PREP Self-Assessment Go to details	01/01/2017	12/31/2019	Journal or Self-Assessment	12/31/2019	Launch
Acute Pain Management: Changes and Challenges Go to details	09/27/2016	09/26/2019	Online Course	09/26/2019	Launch
Chronic Pain and the Opioid Crisis Go to details	09/28/2017	09/27/2020	Online Course	09/27/2020	Launch
EQIPP: Bright Futures - Middle Childhood and Adolescence Go to details	04/10/2017	04/09/2020	Online Course	04/09/2020	Launch

Conclusions

- Need more resources for:
 - Mental health
 - Behavioral treatment
 - Alternative medications
 - Multimodal with non-addictive meds
 - Acknowledgement that addiction is a disease
 - Research-treatment, medications, approaches, techniques
 - Education

DO not Need

- MORE ONE SIZE FITS ALL LAWS
- Lawmakers practicing medicine and limiting access to medications
- Insurance companies making poorly thought out rules

Concluding Remarks

- Children are also very responsive to distraction
- Parents can be invaluable
- Assessment is difficult so it is more humane to assume pain
- Treatment and side effects similar to adults



Conclusion

- Peripheral nerve blocks and one shot caudal blocks are fast and simple
- They provide effective pain relief with minimal side effects
- Always calculate toxic LA dose



