

SHORT COMMUNICATION

Seeing gray through the ON and OFF pathways

E.J. CHICHILNISKY¹ AND B.A. WANDELL²

¹Department of Neurobiology, Stanford University, Stanford

²Neuroscience Program and Department of Psychology, Stanford University, Stanford

(RECEIVED June 20, 1995; ACCEPTED October 25, 1995)

Abstract

Color appearance judgments revealed fundamental differences in visual processing of incremental and decremental lights. First, the balance of cone activation required for a light to appear achromatic was different for increments and decrements (Judd, 1940; Helson & Michels, 1948). Second, adaptation—the visual system's adjustment to background light—affected achromatic decrements more than increments. Third, the regulation of adaptation for incremental and decremental stimuli depended differently on background signals from the three cone types. We interpret these asymmetries as differences in mechanisms of adaptation in the ON and OFF pathways, and suggest that they evolved to accommodate the range and physical sources of color signals in the two pathways.

Keywords: Color appearance, Adaptation, Achromatic, Increment, Decrement

Introduction

A prominent feature of the visual system is the segregation of retinal signals into two neural pathways in which neurons are excited mainly by increments (ON pathway) or decrements (OFF pathway) from the background light intensity. In addition to the early observations of Helson and Michels, and Judd (Judd, 1940; Helson & Michels, 1948), in recent years more psychophysical evidence has accumulated for differential processing of increments and decrements (Walraven, 1977; Krauskopf, 1980; White et al., 1980; Whittle, 1986; Bowen et al., 1989; du Buf, 1992; Mausfeld & Niederee, 1993). These have been complemented by physiological measurements suggesting that the ON and OFF pathways mediate detection of incremental and decremental lights (Schiller et al., 1986).

To understand how the visual pathways process increments and decrements to compute color appearance, we made parametric measurements of appearance across a range of adapting conditions. These data revealed systematic differences between increments and decrements in (1) how appearance is computed, and (2) how appearance varies with adaptation. We interpret these asymmetries as differences in the mechanisms of adaptation in the ON and OFF pathways.

Methods

Overview

To analyze the role of incremental and decremental cone photoreceptor signals in color appearance, we estimated (see below)

the cone absorptions required to cancel the hue of a stimulus. Targets were presented on a large uniform background whose color was set by the experimenter. After adapting to the background light for 2 min, subjects adjusted the spectral composition of the target to make it appear achromatic. Subjects made achromatic settings easily and consistently for a range of targets lighter and darker than the background, presented in random order, yielding achromatic shades ranging from black to gray to white. These settings establish the combination of L(ong), M(iddle), and S(hort) wavelength-sensitive cone absorptions needed to achieve achromatic appearance. Hence we measure color appearance by cancelling it.

Stimuli

Stimuli were presented on a color computer monitor (Hitachi HM-4320D) driven by a NNGS video card from an IBM PC. Targets were 1.5-deg squares viewed foveally at the center of a 45 deg (vertical) × 60 deg (horizontal) background. The visual field beyond the monitor was filled with a reflective viewing hood (75% reflectance for all wavelengths between 370 and 730 nm).

The monitor was calibrated for phosphor independence and nonlinearity using standard methods (Brainard, 1989; Wandell, 1995). To estimate the cone quantum absorptions, phosphor spectral power distributions ($\mu\text{W}/\text{cm}^2\text{-nm-sr}$, PhotoResearch PR-703A) were multiplied by the human cone spectral sensitivity estimates of Smith and Pokorny (Smith & Pokorny, 1975). Each cone spectral sensitivity was normalized to a peak of one.

Subjects set matches on a total of 387 backgrounds. These backgrounds spanned most of the range of luminances and chromaticities that could be achieved on the monitor. The background luminances (CIE 1931) ranged from about 20 to

Reprint requests to: E.J. Chichilnisky, Department of Neurobiology, Stanford University, Stanford, CA, 94305, USA. Internet: ej@white.stanford.edu




Fig. 1. Increment and decrement achromatic settings in two subjects. A: Each point represents the L and S cone quantum absorptions generated by a target adjusted by the observer to appear achromatic, expressed as differences (ΔL , ΔS) from the background. ΔM values, which are approximately proportional to ΔL values in these data, are omitted for simplicity. Increment and decrement cone absorption ratios $\Delta S/\Delta L$ were estimated separately and are indicated by solid lines. Dashed lines extend the solid lines into the opposite quadrant to demonstrate that increment and decrement cone absorption ratios differ. Subject: SY, Background chromaticity (0.29, 0.41; CIE, 1931). B: Similar to A, but for subject RR. Background chromaticity: (0.22, 0.19).

80 cd/m². The background chromaticities covered a circular region of the CIE 1931 (x , y) chromaticity diagram centered at (0.3, 0.3) with diameter 0.2.* As with most monitors, this range of chromaticity coordinates does not approach the spectrum locus.

Subjects

Subjects adjusted the target to appear achromatic by varying the intensity of each of the three monitor phosphors at the target location using a button box. Three paid undergraduates served as subjects (female subjects SY and LW, male subject RR). All subjects had normal color vision according to the Ishihara plates (Ishihara, 1977).

Results

Achromatic appearance

Fig. 1A shows the estimated cone absorptions produced by a collection of achromatic targets.† The coordinates of each point show the effect of one achromatic target on the L and S cones, expressed as differences (ΔL , ΔS) from the absorptions due to the background light, whose intensity was fixed. (For simplicity, the effect on the M cones is not shown.) We refer to the

stimuli in the all-positive quadrant as increments because they excite all three cone types more than the background light. Decremental stimuli excite the cones less than the background light and are plotted in the all-negative quadrant. Increments include light grays and white, and decrements include dark grays and black.

Achromatic appearance is computed differently for incremental and decremental cone signals. The solid line through the increments shows where these data would fall if the visual system classified as achromatic those lights that generate a particular ratio of cone absorptions ($\Delta S/\Delta L = 0.83 \pm 0.06$). This type of model fits increments well (Werner & Walraven, 1982). However, this ratio fails to make decrements appear achromatic. This is shown by the dashed continuation of the increment line into the opposite quadrant, which systematically misses the decrement settings. Instead, decrements appear achromatic when they generate a smaller ratio of cone absorptions ($\Delta S/\Delta L = 0.39 \pm 0.01$), indicated by the solid line in the negative quadrant. Fig. 1B shows similar data for another subject, measured on a different background. In this case, the increment $\Delta S/\Delta L$ ratio required for achromatic appearance was smaller than the decrement ratio. Which ratio was larger varied with the background and the subject. In 6335 achromatic settings made by three subjects, increment and decrement $\Delta S/\Delta L$ ratios differed ($P < 0.01$) for 127 out of 387 backgrounds tested. These systematic differences suggest that two different pathways (e.g. the ON and OFF pathways) process incremental and decremental cone signals.

Light adaptation

Light adaptation, the visual system's adjustment to background light, affected increment and decrement achromatic appearance asymmetrically. This was shown by examining how subjects'

*A complete list of the background luminance and chromaticities may be obtained on the internet at ej@white.stanford.edu

†Target color appearance in these conditions is probably due to signals initiated in the cones because (a) there are few rods in the central 1.5 deg of the retina, and (b) targets were presented on high mesopic or low photopic backgrounds (Wyszecki & Stiles, 1982, p. 406). We will discuss the possibility of rod contributions to adaptation in the Discussion.

achromatic judgments varied with the background light. Fig. 2A shows cone absorptions for achromatic targets seen on two backgrounds which differ only in their S-cone stimulation. Filled symbols represent targets seen on one background, and open symbols represent targets seen on the other. The latter have a higher slope: a larger $\Delta S/\Delta L$ ratio is needed to achieve achromatic appearance on the second background, which stimulated the S-cones more. However, increment ratios change differently than decrement ratios. For example, the background change of Fig. 2A changed the $\Delta S/\Delta L$ increment cone absorption ratio by a factor of 2.65 ± 0.29 , but changed the decrement ratio by a factor of 6.01 ± 0.18 . This is reflected in the larger slope difference between the two decrement lines than the two increment lines.

As the background light changed, the appearance of increments and decrements each varied systematically but according to different rules. Fig. 3A shows how the $\Delta S/\Delta L$ ratio for incremental and decremental achromatic targets varied on backgrounds that differed only in their S-cone stimulation. Filled circles show the ratios for decremental achromatic settings, and open symbols show the ratios for incremental settings. These have been normalized by the ratio on the first background to simplify the visual comparison. Although $\Delta S/\Delta L$ ratios for the increments and decrements both increased with S cone background light, and latter increased linearly and substantially more.

Changes in M and L cone background absorptions also revealed substantial differences between increment and decrement adaptation. These results are shown in Figs. 2B–2C and 3B–3C. Changes in background light always affected decrements more than increments. Large systematic differences in adaptation for increments and decrements, like those shown in Fig. 3, were also observed in two other subjects.

How do background absorptions in one cone type affect sensitivity to increment and decrement signals in the other cone types? The effect of background light on decrement appearance was consistent with independent adaptation of signals from each cone type. The filled squares in Fig. 3D show that varying the S cone background absorptions did not affect the L and M absorptions required to make decrements appear achromatic. Similarly, varying either the M or L cone background absorptions did not affect the cone absorptions of achromatic decrement measurements in the other two cone classes (filled circles in Figs. 3E–3F). The fact that adaptation affects achromatic decrement appearance independently in each cone class suggests that segregation into incremental and decremental pathways takes place prior to neural color opponency.

Again, adaptation measured by increments followed different rules. Varying the background absorptions in one cone type affected the cone absorptions in the other two cone classes required to make increments appear achromatic (open circles in Figs. 3D–3F) (Werner & Walraven, 1982). The effect of adaptation reversed as the background hue passed to its complement through neutral gray. This was observed in all three subjects. These data suggest that neural opponent mechanisms affect sensitivity to incremental cone signals (Pugh, 1976; Mollon & Polden, 1977; Sternheim et al., 1979; Wandell & Pugh, 1980a).

Discussion

Our results may be summarized as follows. First, the cone absorption ratios of achromatic increment and decrement lights differed. Second, these ratios were affected differently by adaptation. What do these findings suggest about the organization of the visual pathways?




Fig. 2. Effect of background changes on achromatic settings. A: Achromatic settings gathered on two backgrounds that differed only in their effect on the S cones. Stimuli are represented as differences (ΔL , ΔS) from their respective backgrounds. Open symbols represent stimuli seen on the first background (8.76 units of S cone absorption), and filled symbols represent stimuli seen on the second background (1.38 units of S cone absorption). The background change caused the estimated increment ratio $\Delta S/\Delta L$ to change by a factor of 2.65 ± 0.29 , and the decrement ratio by a factor of 6.01 ± 0.18 . Ratios were computed using multiple regression; “ \pm ” indicates a 95% confidence interval computed by separately resampling increment and decrement data 250 times (Efron, 1982). Other conditions are as in Fig. 1. B: ΔL and ΔM absorptions for two data sets gathered on backgrounds that differ only in their effect on the M cones. Open symbols show data from a background with 8.81 units of M cone absorption, and filled symbols from a background with 7.08 units of M cone absorption. The background change caused the increment $\Delta M/\Delta L$ ratio to change by a factor of 1.12 ± 0.02 , decrement ratio by a factor of 1.22 ± 0.01 . C: ΔM and ΔL absorptions for two data sets gathered on backgrounds that differ only in their effect on the L cones. Open symbols show data from a background with 8.92 units of L cone absorption, and filled symbols from a background with 7.19 units of L cone absorption. The background change caused the increment $\Delta L/\Delta M$ ratio to change by a factor of 1.12 ± 0.02 , decrement ratio by a factor of 1.23 ± 0.01 . Subject: RR.




Fig. 3. Effect of many background changes on achromatic increment and decrement absorption ratios. A: Increment and decrement $\Delta S/\Delta L$ cone absorption ratios of achromatic settings gathered on seven backgrounds which differed only in their effect on the S cones. The horizontal axis indicates S cone absorptions due to each background. Each curve has been normalized by the value on the first background to simplify visual comparison. B: $\Delta M/\Delta L$ ratios as a function of background changes selective for the M cones. C: $\Delta L/\Delta M$ ratios as a function of background changes selective for the L cones. D: Increment and decrement $\Delta L/\Delta M$ for the same data as A. Labels on horizontal axes indicate appearance of background light in a dark room. E: Increment and decrement $\Delta L/\Delta S$ for the same data as B. F: Increment and decrement $\Delta M/\Delta S$ for the same data as C. Subject: RR.

Primate cone photocurrent varies linearly in response to dim flashes (Schnapf et al., 1990). It is therefore likely that cones respond symmetrically to increments and decrements of moderate contrast around the photopic backgrounds used in our experiments. If this is true, sensitivity changes in cone photocurrent (Schnapf et al., 1990) cannot explain the asymmetries in increment and decrement appearance changes with background light (Figs. 2 and 3).

To manage transduction efficiently, it is sensible that the cones themselves should adapt. Moreover, there is considerable psychophysical and physiological evidence that cone sensitivity depends on the background illumination (Stiles, 1959; Boynton & Whitten, 1970; Schnapf et al., 1990) and that this, in turn, affects subjects' judgments of color appearance (von Kries, 1905; Walters, 1942; Wright, 1947; Burnham et al., 1952; Hurvich & Jameson, 1958; Wassef, 1958; Walraven, 1976; Shevell, 1978; Werner & Walraven, 1982; Brainard & Wandell, 1992; Chichilnisky & Wandell, 1995). There is also a substantial body of psychophysical evidence documenting post-receptor adaptation under certain viewing conditions (Pugh, 1976; Mollon & Polden, 1977; Pugh & Mollon, 1979; Pugh & Larimer, 1980; Wandell & Pugh, 1980a,b; Stromeyer & Sternheim, 1981; Ahn & Macleod, 1993). The results we report here are consistent with a mixture of receptor and postreceptor adaptation. When judging the appearance of decremental lights, subjects' behavior is consistent with separate adaptation within each cone class (Fig. 3, filled circles). When judging the appearance of incremental lights, subjects' behavior is consistent with adaptation being controlled by signals from more than one receptor class (Fig. 3, open circles). A natural explanation is that the anatomically distinct ON and OFF pathways regulate sensitivity according to different rules. For example, in ON-bipolar cells surround signals from one cone type may affect the sensitivity to signals from another cone type; while in OFF-bipolar cells this interaction may be absent.

As mentioned previously, target appearance is probably based on signals initiated in the foveal cones. However, peripheral photoreceptor signals, including rod signals, may contribute to asymmetric processing of increments and decrements by affecting sensitivity of signals initiated in foveal cones. Consider, for example, the asymmetric effect of S cone selective background changes on increment and decrement appearance, shown in Figs. 3A and 3D. These background changes may have also affected rod signals, which could have asymmetric effects on sensitivity of ON and OFF signals initiated by foveal cones. Whether or not the rods play a role, the background changes in Fig. 3D were invisible to the L and M cones, whose ratio is plotted on the vertical axis. Thus, the data demonstrate interactions between receptor types in increment but not decrement adaptation.

Our findings suggest a neural basis for a classical perceptual observation known as the Helson-Judd effect (Judd, 1940; Helson & Michels, 1948). These authors observed that targets that are substantially less intense than the background appear achromatic at chromaticities similar to that of the background; targets that are substantially more intense than the background appear achromatic at a background-independent chromaticity.† Our observations show that there are sharp differences in cone

absorption ratios (Fig. 1) and adaptation (Figs. 2–3) for achromatic increments and decrements of moderate contrast. This suggests that the neural basis of the Helson-Judd effect is asymmetric processing and adaptation in the ON and OFF pathways.

The data in Fig. 3 suggest that more complete adaptation occurs in the OFF pathway, and that light adaptation in the OFF pathway is regulated separately within each cone type whereas adaptation in the ON pathway depends in part on opponent-color signals. Why should there be different rules for adaptation in the ON and OFF pathways? Stimuli that excite these pathways require different visual processing because (1) the range of increment intensities exceeds the range of decrement intensities (increments may be arbitrarily intense while decrements are bounded by zero), and (2) light sources and specular reflections typically create increments, while diffuse reflections typically create decrements. The ON and OFF pathways may adapt differently because these two pathways usually encode stimuli of different physical origins.

Acknowledgments

We thank D. Baylor, S. du Lac, R. Fernald, D. Heeger, and W. Newsome for comments on the manuscript. This work was supported by NEI Grant RO1-EY03164-16 and NASA Grant NCC-2-307. E.J. Chichilnisky was supported by a predoctoral fellowship from the Howard Hughes Medical Institute.

References

- AHN, S.J. & MACLEOD, D.I.A. (1993). Link-specific adaptation in the luminance and chromatic channels. *Vision Research* **33**, 2271–2286.
- BOWEN, R.W., POKORNY, J. & SMITH, V.C. (1989). Sawtooth contrast sensitivity: Decrements have the edge. *Vision Research* **29**, 1501–1509.
- BOYNTON, R.M. & WHITTEN, D.N. (1970). Visual adaptation in monkey cones: Recordings of late receptor potentials. *Science* **170**, 1423–1426.
- BRAINARD, D.H. (1989). Calibration of a computer controlled color monitor. *Color Research and Application* **14**, 23–34.
- BRAINARD, D.H. & WANDELL, B.A. (1992). Asymmetric color-matching: How color appearance depends on the illuminant. *Journal of the Optical Society of America A* **9**, 1433–1448.
- BURNHAM, R.W., EVANS, R.M. & NEWHALL, S.M. (1952). Influence on color perception of adaptation to illumination. *Journal of the Optical Society of America* **42**, 597–605.
- CHICHILNISKY, E.J. & WANDELL, B.A. (1995). Photoreceptor sensitivity changes explain color appearance shifts induced by large uniform backgrounds in dichoptic matching. *Vision Research* **35**, 239–254.
- DU BUF, J.M.H. (1992). Brightness versus apparent contrast: I. Incremental and decremental disks with varying diameter. *Spatial Vision* **6**, 159–182.
- EFRON, B. (1982). *The Jackknife, the Bootstrap and Other Resampling Plans*. Philadelphia, Pennsylvania: Society for Industrial and Applied Mathematics.
- HELSON, H. & MICHELS, W.C. (1948). The effect of chromatic adaptation on achromaticity. *Journal of the Optical Society of America* **38**, 1025–1032.
- HUNT, R.W.G. (1995). *The Reproduction of Color*. Tolworth, England: Fountain Press.
- HURVICH, L.M. & JAMESON, D. (1958). Further development of a quantified opponent-color theory. In *Visual Problems of Colour II*, pp. 691–723. London: HMSO.
- ISHIHARA, S. (1977). *Tests for Colour-Blindness*. Tokyo: Kanehara Shuppen Company, Ltd.
- JUDD, D.B. (1940). Hue saturation and lightness of surface colors with chromatic illumination. *Journal of the Optical Society of America* **30**, 21–32.
- KRAUSKOPF, J. (1980). Discrimination and detection of changes in luminance. *Vision Research* **20**, 671–677.
- MAUSFELD, R. & NIEDEREE, R. (1993). An inquiry into relational con-

†The Helson-Judd effect is sometimes described in terms of illuminants and surfaces; see, for example, Hunt (1995), p. 716.

- cepts of colour, based on incremental principles of colour coding for minimal relational stimuli. *Perception* **22**, 427-462.
- MOLLON, J.D. & POLDEN, P.G. (1977). An anomaly in the response of the eye to light to short wavelengths. *Philosophical Transactions of the Royal Society B* (London) **278**, 207-240.
- PUGH, E.N. (1976). The nature of the π_1 colour mechanism of W.S. Stiles. *Journal of Physiology* (London) **257**, 713-747.
- PUGH, E.N. & LARIMER, J. (1980). Test of the identity of the site of blue/yellow hue cancellation and the site of chromatic antagonism in the π_1 pathway. *Vision Research* **20**, 779-788.
- PUGH, E.N. & MOLLON, J.D. (1979). A theory of the π_1 and π_3 color mechanisms of Stiles. *Vision Research* **19**, 293-312.
- SCHILLER, P.H., SANDELL, J.H. & MAUNSELL, J.H.R. (1986). Functions of the ON and OFF channels of the visual system. *Nature* **322**, 824-825.
- SCHNAPF, J.L., NUNN, B.J., MEISTER, M. & BAYLOR, D.A. (1990). Visual transduction in cones of the monkey *Macaca fascicularis*. *Journal of Physiology* (London) **427**, 681-713.
- SHEVELL, S.K. (1978). The dual role of chromatic backgrounds in color perception. *Vision Research* **18**, 1649-1661.
- SMITH, V. & POKORNY, J. (1975). Spectral sensitivity of the foveal cone photopigments between 400 and 500 nm. *Vision Research* **15**, 161-171.
- STERNHEIM, C.E., STROMEYER, C.F., III & KHOO, M.C.K. (1979). Visibility of chromatic flicker upon spectrally mixed adapting fields. *Vision Research* **19**, 175-183.
- STILES, W.S. (1959). Color vision: The approach through increment threshold sensitivity. *Proceedings National Academy of Sciences of the U.S.A.* **45**, 100-114.
- STROMEYER, C.F., III & STERNHEIM, C.E. (1981). Visibility of red and green spatial patterns upon spectrally mixed adapting fields. *Vision Research* **21**, 397-408.
- VON KRIES, J. (1905). Influence of adaptation on the effects produced by luminous stimuli. In *Sources of Color Science*, ed. MACADAM, D.L., pp. 120-127. Cambridge, Massachusetts: MIT Press.
- WALRAVEN, J. (1976). Discounting the background—the missing link in the explanation of chromatic induction. *Vision Research* **16**, 289-295.
- WALRAVEN, J. (1977). Colour signals from incremental and decremental light stimuli. *Vision Research* **17**, 71-76.
- WALTERS, H.V. (1942). Some experiments on the trichromatic theory of vision. *Proceedings of the Royal Society B* (London) **131**, 27-50.
- WANDELL, B.A. (1995). *Foundations of Vision*. Sunderland, Massachusetts: Sinauer.
- WANDELL, B.A. & PUGH, E.N. (1980a). Detection of long-duration, long-wavelength incremental flashes by a chromatically coded pathway. *Vision Research* **20**, 625-636.
- WANDELL, B.A. & PUGH, E.N. (1980b). A field-additive pathway detects brief-duration, long-wavelength incremental flashes. *Vision Research* **20**, 613-624.
- WASSEF, E.G.T. (1958). Investigation into the theory of prediction of the appearance of colors and its bearing on the theory of color vision. *Optica Acta* **5**, 101-108.
- WERNER, J.S. & WALRAVEN, J. (1982). Effect of chromatic adaptation on the achromatic locus: The role of contrast, luminance and background color. *Vision Research* **22**, 929-944.
- WHITE, T.W., IRVIN, G.E. & WILLIAMS, M.C. (1980). Asymmetry in the brightness and darkness Broca-Sulzer effects. *Vision Research* **20**, 723-726.
- WHITTLE, P. (1986). Increments and decrements: Luminance discrimination. *Vision Research* **26**, 1677-1692.
- WRIGHT, W.D. (1947). *Researches on Normal and Defective Colour Vision*. St. Louis, Missouri: C.V. Mosby.
- WYSZECKI, G. & STILES, W.S. (1982). *Color Science—Concepts and Methods, Quantitative Data and Formulae*. New York: John Wiley & Sons.